

Topological interpretations of provability logic

Lev Beklemishev*

David Gabelaia†

October 30, 2012

Abstract

Provability logic concerns the study of modality \Box as provability in formal systems such as Peano arithmetic. Natural, albeit quite surprising, topological interpretation of provability logic has been found in the 1970's by Harold Simmons and Leo Esakia. They have observed that the dual \Diamond modality, corresponding to consistency in the context of formal arithmetic, has all the basic properties of the topological derivative operator acting on a scattered space. The topic has become a long-term project for the Georgian school of logic led by Esakia, with occasional contributions from elsewhere.

More recently, a new impetus came from the study of polymodal provability logic **GLP** that was known to be Kripke incomplete and, in general, to have a more complicated behavior than its unimodal counterpart. Topological semantics provided a better alternative to Kripke models in the sense that **GLP** was shown to be topologically complete. At the same time, new fascinating connections with set theory and large cardinals have emerged.

We give a survey of the results on topological semantics of provability logic starting from first contributions by Esakia. However, a special emphasis is put on the recent work on topological models of polymodal provability logic. We also included a few results that have not been published so far, most notably the results of Section 6 (due the second author) and Sections 10, 11 (due to the first author).

1 Provability algebras and logics

Provability logics and algebras emerge from, respectively, a modal logical and an algebraic point of view on the proof-theoretic phenomena around Gödel's incompleteness theorems. These theorems are usually perceived as putting fundamental restrictions on what can be formally proved in a given axiomatic system (satisfying modest natural requirements). For the sake of a discussion, we call a formal theory T *gödelian* if

- T is a first order theory in which the natural numbers along with the operations $+$ and \cdot are interpretable;

*V.A. Steklov Mathematical Institute, RAS; Moscow M.V. Lomonosov State University; National Research University Higher School of Economics; email: bekl@mi.ras.ru

†TSU Razmadze Institute of Mathematics, Tbilisi; email: gabelaia@gmail.com

- T proves some basic properties of these operations and a modicum of induction (it is sufficient to assume that T contains the Elementary Arithmetic EA);
- T has a recursively enumerable set of axioms.

The Second Incompleteness Theorem of Kurt Gödel (G2) states that a gödelian theory T cannot prove its own consistency provided it is indeed consistent. More accurately, for any r.e. presentation of such a theory T , Gödel has shown how to write down an arithmetical formula $\text{Prov}_T(x)$ expressing that x is (a natural number coding) a formula provable in T . Then the statement $\text{Con}(T) := \neg \text{Prov}_T(\ulcorner \perp \urcorner)$ naturally expresses that the theory T is consistent. G2 states that $T \not\vdash \text{Con}(T)$ provided T is consistent.

Provability logic emerged from the question what properties of formal provability Prov_T can be verified in T , even if the consistency of T cannot. Several such properties have been stated by Gödel himself [30]. Hilbert and Bernays and then Löb [38] stated them in the form of conditions any adequate formalization of a provability predicate in T must satisfy. After Gödel's and Löb's work it was clear that the formal provability predicate calls for a treatment as a modality. It led to the formulation of the Gödel–Löb provability logic **GL** and eventually to the celebrated arithmetical completeness theorem due to Robert Solovay.

Independently, Macintyre and Simmons [39] and Magari [40] took a very natural algebraic perspective on the phenomenon of formal provability which led to the concept of *diagonalizable algebra*. Such algebras are now more commonly called *Magari algebras*. This point of view will be more convenient for our present purposes.

Recall that the Lindenbaum–Tarski algebra of a theory T is the set of all T -sentences Sent_T modulo provable equivalence in T , that is, the structure $\mathcal{L}_T = \text{Sent}_T / \sim_T$ where, for all $\varphi, \psi \in \text{Sent}_T$,

$$\varphi \sim_T \psi \iff T \vdash (\varphi \leftrightarrow \psi).$$

Since we assume T to be based on classical propositional logic, \mathcal{L}_T is a boolean algebra with operations \wedge, \vee, \neg . Constants \perp and \top are identified with the sets of refutable and of provable sentences of T , respectively. The standard ordering on \mathcal{L}_T is defined by

$$[\varphi] \leq [\psi] \iff T \vdash \varphi \rightarrow \psi \iff [\varphi \wedge \psi] = [\varphi],$$

where $[\varphi]$ denotes the equivalence class of φ .

It is well-known that for consistent gödelian T all such algebras are isomorphic to the unique countable atomless boolean algebra. (This is a consequence of a strengthening of Gödel's First Incompleteness Theorem due to Rosser.) We obtain more interesting algebras by enriching the structure of the boolean algebra \mathcal{L}_T by additional operation(s).

Gödel's consistency formula induces a unary operator \Diamond_T acting on \mathcal{L}_T :

$$\Diamond_T : [\varphi] \mapsto [\text{Con}(T + \varphi)].$$

The sentence $\text{Con}(T + \varphi)$ expressing the consistency of T extended by φ can be defined as $\neg \text{Prov}_T(\ulcorner \neg \varphi \urcorner)$. The dual operator is $\Box_T : [\varphi] \mapsto [\text{Prov}_T(\ulcorner \varphi \urcorner)]$, thus $\Box_T x = \neg \Diamond_T \neg x$, for all $x \in \mathcal{L}_T$.

Bernays–Löb derivability conditions ensure that \Diamond_T is correctly defined on the equivalence classes of the Lindenbaum–Tarski algebra of T . Moreover, it satisfies the following identities (where we write \Diamond_T simply as \Diamond and the variables range over arbitrary elements of \mathcal{L}_T):

$$\text{M1. } \Diamond \perp = \perp; \quad \Diamond(x \vee y) = \Diamond x \vee \Diamond y;$$

$$\text{M2. } \Diamond x = \Diamond(x \wedge \neg \Diamond x).$$

Notice that Axiom M2 is a formalization of G2 stated for the theory $T' = T + \varphi$, where $[\varphi] = x$. In fact, the left hand side states that T' is consistent, whereas the right hand side states that $T' + \neg \text{Con}(T')$ is consistent, that is, $T' \not\vdash \text{Con}(T')$. The dual form of Axiom M2,

$$\Box(\Box x \rightarrow x) = \Box x,$$

expresses the familiar Löb’s theorem.

Notice also that M1 induces \Diamond to be monotone: if $x \leq y$ then $\Diamond x \leq \Diamond y$.

Boolean algebras with operators $\mathcal{M} = (M, \Diamond)$ satisfying the above identities are called *Magari algebras*. Thus, the main example of a Magari algebra is the structure $(\mathcal{L}_T, \Diamond_T)$, for any consistent gödelian theory T , which is also called the *provability algebra of T* .

The *transitivity* inequality $\Diamond \Diamond x \leq \Diamond x$ is often postulated as an additional axiom of Magari algebras, however it follows from M1 and M2.

Proposition 1.1. *In any Magari algebra \mathcal{M} , there holds $\Diamond \Diamond x \leq \Diamond x$, for all $x \in M$.*

Proof. Given any $x \in M$ consider $y := x \vee \Diamond x$. On the one hand, we have

$$\Diamond \Diamond x \leq (\Diamond x \vee \Diamond \Diamond x) = \Diamond y.$$

On the other hand, since $\Diamond x \wedge \neg \Diamond y = \perp$ we obtain

$$\Diamond y \leq \Diamond(y \wedge \neg \Diamond y) \leq \Diamond((x \vee \Diamond x) \wedge \neg \Diamond y) \leq \Diamond(x \wedge \neg \Diamond y) \vee \Diamond \perp \leq \Diamond x.$$

Hence, $\Diamond \Diamond x \leq \Diamond x$. ◻

In general, we call an *identity* of an algebraic structure \mathcal{M} a formula of the form $t(\vec{x}) = u(\vec{x})$, where t, u are terms, such that $\mathcal{M} \models \forall \vec{x} (t(\vec{x}) = u(\vec{x}))$. Identities of Maragi algebras can be described in terms of modal logic as follows. Any term (built from the variables using boolean operations and \Diamond) is naturally identified with a formula in the language of propositional logic with a new unary connective \Diamond . If $\varphi(\vec{x})$ is such a formula and \mathcal{M} a Magari algebra, we write $\mathcal{M} \models \varphi$ iff $\forall \vec{x} (t_\varphi(\vec{x}) = \top)$ is valid in \mathcal{M} , where t_φ is the term corresponding to φ . Since any identity in Magari algebras can be equivalently written in the form $t = \top$, for some term t , the axiomatization of identities of \mathcal{M} amounts to axiomatizing modal formulas valid in \mathcal{M} . The *logic of \mathcal{M}* , $\text{Log}(\mathcal{M})$, is the set of all modal formulas valid in \mathcal{M} , that is,

$$\text{Log}(\mathcal{M}) := \{\varphi : \mathcal{M} \models \varphi\},$$

and the logic of a class of modal algebras is defined similarly.

One of the main parameters of a Magari algebra \mathcal{M} is its *characteristic* $\text{ch}(\mathcal{M})$ which is defined as follows. We let

$$\text{ch}(\mathcal{M}) := \min\{k \in \omega : \Diamond^k \top = \perp\}$$

and $\text{ch}(\mathcal{M}) := \infty$, if no such k exists. If T is arithmetically sound, that is, if the arithmetical consequences of T are valid in the standard model, then $\text{ch}(\mathcal{L}_T) = \infty$. Theories of finite characteristics are, in a sense, close to being inconsistent and can be considered a pathology.

Robert Solovay in his fundamental paper [46] proved that any identity valid in the structure $(\mathcal{L}_T, \Diamond_T)$ follows from the boolean identities together with M1–M2, provided T is arithmetically sound. This has been generalized by Albert Visser [48] to arbitrary theories of infinite characteristic. Put algebraically, this becomes the following statement.

Theorem 1.2. *Suppose $\text{ch}(\mathcal{L}_T, \Diamond_T) = \infty$. An identity holds in $(\mathcal{L}_T, \Diamond_T)$ iff it holds in all Magari algebras.*

Apart from the equational characterization by M1–M2 above, the identities of Magari algebras can be axiomatized modal-logically. In fact, the logic of all Magari algebras, and by Solovay theorem the logic $\text{Log}(\mathcal{L}_T, \Diamond_T)$ of the Magari algebra of T , for any fixed theory T of infinite characteristic, coincides with the familiar Gödel–Löb logic **GL**. By abuse of language we will often identify **GL** with the set of identities of Magari algebras.

A Hilbert-style axiomatization of **GL** is usually given in the modal language where \Box rather than \Diamond is taken as basic and the latter is treated as an abbreviation for $\neg\Box\neg$. The axioms and inference rules of **GL** are as follows.

Axiom schemata:

- L1. All instances of propositional tautologies;
- L2. $\Box(\varphi \rightarrow \psi) \rightarrow (\Box\varphi \rightarrow \Box\psi)$;
- L3. $\Box(\Box\varphi \rightarrow \varphi) \rightarrow \Box\varphi$.

Rules: $\varphi, \varphi \rightarrow \psi / \psi$ (modus ponens), $\varphi / \Box\varphi$ (necessitation).

By a well-known result of Segerberg, **GL** is sound and complete w.r.t. the class of all transitive and upwards well-founded Kripke frames. In fact, it is sufficient to restrict the attention to frames that are finite irreflexive trees. Thus, summarizing various characterizations above, we have

Theorem 1.3. *For any modal formula φ , the following statements are equivalent:*

- (i) **GL** $\vdash \varphi$;
- (ii) φ is valid in all Magari algebras;
- (iii) $(\mathcal{L}_T, \Diamond_T) \models \varphi$, for any T of infinite characteristic;
- (iv) φ is valid in all finite irreflexive tree-like Kripke frames.

2 Topological interpretation

Having defined the notion of Magari algebra, the first question one can ask is whether there are any other natural examples of such algebras apart from the provability algebras of the form $(\mathcal{L}_T, \Diamond_T)$. The fact that such algebras naturally emerge from scattered topological spaces was discovered independently by Harold

Simmons [44] and Leo Esakia [27].¹ We now infer this semantics from rather general considerations.

It is a standard idea in logic, going back at least as early as to Leibnitz, to interpret propositions as subsets of a given nonempty set X . Then, logical connectives are naturally translated as the familiar boolean operations on sets. From this point of view, \Diamond should correspond to an operator acting on the subsets of X , so we come to the following mathematical question.

Let X be a nonempty set, $\mathcal{P}(X)$ the boolean algebra of subsets of X . Consider any operator $\delta : \mathcal{P}(X) \rightarrow \mathcal{P}(X)$ and the structure $(\mathcal{P}(X), \delta)$. Can $(\mathcal{P}(X), \delta)$ be a Magari algebra and, if yes, when? Simmons and Esakia, in some sense, found a canonical answer to this question.

Let (X, τ) be a topological space, where τ denotes the set of open subsets of X , and let $A \subseteq X$. Topological *derivative* $d_\tau(A)$ of A is the set of limit points of A :

$$x \in d_\tau(A) \iff \forall U \in \tau (x \in U \Rightarrow \exists y \neq x (y \in U \cap A)).$$

Notice that $c_\tau(A) := A \cup d_\tau(A)$ is the closure of A and $i_\tau(A) := A \setminus d_\tau(A)$ is the set of isolated points of A .

The classical notion of a scattered topological space is due to Georg Cantor. (X, τ) is called *scattered* if every nonempty subspace $A \subseteq X$ has an isolated point.

Theorem 2.1 (Simmons, Esakia). *The following statements are equivalent:*

- (i) (X, τ) is scattered;
- (ii) For all $A \subseteq X$, $d_\tau(A) = d_\tau(A \setminus d_\tau(A))$;
- (iii) $(\mathcal{P}(X), d_\tau)$ is a Magari algebra.

Notice that (ii) means that each point of A is a limit point of its isolated points. Algebras of the form $(\mathcal{P}(X), d_\tau)$ associated with a topological space (X, τ) will be called *derivative algebras*. Thus, this theorem states that the derivative algebra of (X, τ) is Magari iff (X, τ) is scattered.

Proof. First, we notice that Statement (iii) is just a reformulation of (ii). In fact, the identities $d_\tau \emptyset = \emptyset$ and $d_\tau(A \cup B) = d_\tau(A) \cup d_\tau(B)$ are obviously valid in all topological spaces. We show the equivalence of (i) and (ii).

Suppose (X, τ) is scattered, $A \subseteq X$ and $x \in d_\tau(A)$. Consider any open neighborhood U of x . Since $U \cap A \setminus \{x\}$ is nonempty, it has an isolated point $y \neq x$. Since U is open, y is an isolated point of A , that is, $y \in A \setminus d_\tau(A)$. Hence, $x \in d_\tau(A \setminus d_\tau(A))$. The inclusion $d_\tau(A \setminus d_\tau(A)) \subseteq d_\tau(A)$ follows from the monotonicity of d_τ . Therefore Statement (ii) holds.

Suppose that (ii) holds and let $A \subseteq X$ be nonempty. We show that A has an isolated point. If $d_\tau A$ is empty, we are done. Otherwise, take any $x \in d_\tau A$. Since x is a limit of isolated points of A , there must be at least one such point. \dashv

We notice that the transitivity principle $d_\tau d_\tau A \subseteq d_\tau A$ topologically means that the set $d_\tau A$, for any $A \subseteq X$, is closed. We recall the following standard equivalent characterization an easy proof of which we shall omit.

¹The paper by Simmons was published in the pre-Solovay era of provability logic and remained largely ignored. Esakia apparently independently discovered the interpretation when he was already familiar with the works of Magari and Solovay.

Proposition 2.2. *For any topological space (X, τ) , the following statements are equivalent:*

- (i) *Every $x \in X$ is an intersection of an open and a closed set;*
- (ii) *For each $A \subseteq X$, the set $d_\tau A$ is closed.*

Topological spaces satisfying either of these conditions are called T_d -spaces. Condition (i) shows that T_d is a weak separation property located between T_0 and T_1 . Thus, Proposition 1.1 yields the following corollary.

Corollary 2.3. *All scattered spaces are T_d .*

We have seen in Theorem 2.1 that each scattered space equipped with a topological derivative operator is a Magari algebra. The following result shows that any Magari algebra on $\mathcal{P}(X)$ can be described in this way.

Theorem 2.4. *If $(\mathcal{P}(X), \delta)$ is a Magari algebra, then X bears a unique topology τ for which $\delta = d_\tau$. Moreover, τ is scattered.*

Proof. We first remark that, if $(\mathcal{P}(X), \delta)$ is a Magari algebra, then the operator $c(A) := A \cup \delta A$ satisfies the Kuratowski axioms of the topological closure: $c\emptyset = \emptyset$, $c(A \cup B) = cA \cup cB$, $A \subseteq cA$, $ccA = cA$ (the latter uses the transitivity property of δ). This defines a topology τ on X in which a set A is closed iff $A = c(A)$ iff $\delta A \subseteq A$. If δ coincides with d_τ , this condition is also clearly necessary. So, the required topology is unique. To show that $\delta = d_\tau$ indeed, we need an auxiliary lemma.

Lemma 2.5. *Suppose $(\mathcal{P}(X), \delta)$ is Magari. Then, for all $x \in X$,*

- (i) $x \notin \delta(\{x\})$;
- (ii) $x \in \delta A \iff x \in \delta(A \setminus \{x\})$.

Proof. (i) By Axiom M2 we have $\delta\{x\} \subseteq \delta(\{x\} \setminus \delta\{x\})$. If $x \in \delta\{x\}$ then $\delta(\{x\} \setminus \delta\{x\}) = \delta\emptyset = \emptyset$. Hence, $\delta\{x\} = \emptyset$, a contradiction.

(ii) $x \in \delta A$ implies $x \in \delta((A \setminus \{x\}) \cup \{x\}) = \delta(A \setminus \{x\}) \cup \delta\{x\}$. By (i), $x \notin \delta\{x\}$, hence $x \in \delta(A \setminus \{x\})$. The other implication follows from the monotonicity of δ . \dashv

Lemma 2.6. *Suppose $(\mathcal{P}(X), \delta)$ is Magari and τ is the associated topology. Then $\delta = d_\tau$.*

Proof. Let $d = d_\tau$; we show that, for any set $A \subseteq X$, $dA = \delta A$. Notice that, for any B , $cB = dB \cup B = \delta B \cup B$.

Assume $x \in \delta A$ then

$$x \in \delta(A \setminus \{x\}) \subseteq c(A \setminus \{x\}) \subseteq d(A \setminus \{x\}) \cup (A \setminus \{x\}).$$

Since $x \notin A \setminus \{x\}$, we obtain $x \in d(A \setminus \{x\})$. By the monotonicity of d , $x \in dA$. Similarly, if $x \in dA$ then $x \in d(A \setminus \{x\})$. Hence,

$$x \in c(A \setminus \{x\}) = \delta(A \setminus \{x\}) \cup (A \setminus \{x\}).$$

Since $x \notin A \setminus \{x\}$ we obtain $x \in \delta A$. \dashv

From this lemma and Theorem 2.1 we infer that the associated topology τ must be scattered. \dashv

Theorem 2.4 shows that to study a natural set-theoretic interpretation of provability logic means to study the semantics of \Diamond as a derivative operation on a scattered topological space.

Derivative semantics of modality was first suggested in the fundamental paper by McKinsey and Tarski [42]. See the paper by Kudinov and Shehtman in this volume for a detailed survey of such semantics for general topological spaces. The emphasis in this paper is on the logics related to formal provability. Specifically, scattered spaces (and their generalizations) will play a major role here.

3 Topological completeness theorems

Natural examples of scattered topological spaces come from orderings. Two examples will play an important role below.

Let (X, \prec) be a strict partial ordering. The *left topology* τ_{\leftarrow} on (X, \prec) is given by all sets $A \subseteq X$ such that $\forall x, y (y \prec x \in A \Rightarrow y \in A)$. We obviously have that (X, \prec) is well-founded iff (X, τ_{\leftarrow}) is scattered. The *right topology* or the *upset topology* is defined similarly.

This topology is, in general, non-Hausdorff. More natural is the *order topology* on a linear ordering $(X, <)$, which is generated by all intervals (α, β) such that $\alpha, \beta \in X \cup \{\pm\infty\}$ and $\alpha < \beta$. The order topology refines the left topology and is scattered on any ordinal.

Given a topological space (X, τ) , we shall denote the logic of its derivative algebra $(\mathcal{P}(X), d_\tau)$ by $\text{Log}(X, \tau)$, and we use similar notation for the logic of a class of derivative algebras.

The logic $\text{Log}(X, \tau)$ is often equivalently defined in terms of valuations as follows. A *valuation* on (X, τ) is any map v from the set of propositional formulas to the powerset of X satisfying the following conditions:

- $v(\varphi \vee \psi) = v(\varphi) \cup v(\psi)$, $v(\neg\varphi) = X \setminus v(\varphi)$, $v(\top) = X$, $v(\perp) = \emptyset$,
- $v(\Diamond\varphi) = d(v(\varphi))$.

One usually writes $X, x \models_v \varphi$ for $x \in v(\varphi)$. A formula φ is *valid in X* (denoted $X \models \varphi$) if $\forall v, v(\varphi) = X$. Then, $\text{Log}(X, \tau)$ coincides with the set of all formulas valid in (X, τ) . Similarly, if \mathcal{C} is a class of spaces, $\text{Log}(\mathcal{C})$ is the set of formulas valid in all members $(X, \tau) \in \mathcal{C}$.

Since $\text{Log}(\mathcal{C})$ is the logic of a class of Magari algebras, this set of formulas is closed under the rules modus ponens, necessitation and substitution, and contains all theorems of Gödel–Löb logic **GL**. Thus, $\text{Log}(\mathcal{C})$ is a normal modal logic extending **GL**, for any class \mathcal{C} of scattered spaces.

Topological interpretation of provability logic suggests a question whether **GL** is complete for any natural class of scattered topological spaces. Leo Esakia [27] has shown that **GL** coincides with the logic of the class of all scattered topological spaces. In fact, it is sufficient to take finite trees (considered as strict partial orderings) with the associated upset topology.

Theorem 3.1 (Esakia). (i) $\text{Log}(\mathcal{C}) = \mathbf{GL}$, where \mathcal{C} is the class of all finite irreflexive trees taken with the upset topology.

(ii) *There is a countable scattered space (X, τ) such that $\text{Log}(X, \tau) = \mathbf{GL}$.*

We notice that this theorem follows almost immediately from the completeness of \mathbf{GL} w.r.t. its Kripke semantics. In fact, if (X, \prec) is a strict partial ordering, then the modal algebra associated with the Kripke frame (X, \prec) is the same as the derivative algebra of (X, τ) where τ is its upset topology. This implies that any modal logic of a class of strict partial orders is complete w.r.t. topological semantics.

Esakia's theorem has been strengthened by Abashidze [1] and Blass [18].

Theorem 3.2 (Abashidze, Blass). *Consider an ordinal $\Omega \geq \omega^\omega$ equipped with the order topology. Then $\text{Log}(\Omega) = \mathbf{GL}$.*

Thus, \mathbf{GL} is complete w.r.t. a natural scattered topological space. In Section 5 we give an alternative proof of this result by utilizing the recursive constructions of finite irreflexive trees and ordinals below ω^ω .

4 Some useful techniques

4.1 Ranks and d -maps

An equivalent characterization of scattered spaces is often given in terms of the following transfinite *Cantor–Bendixson sequence* of subsets of a topological space (X, τ) :

- $d_\tau^0 X = X$; $d_\tau^{\alpha+1} X = d_\tau(d_\tau^\alpha X)$ and
- $d_\tau^\alpha X = \bigcap_{\beta < \alpha} d_\tau^\beta X$ if α is a limit ordinal.

It is easy to show by transfinite induction that, for any (X, τ) , all sets $d_\tau^\alpha X$ are closed and that $d_\tau^\alpha X \supseteq d_\tau^\beta X$ whenever $\alpha \leq \beta$.

Theorem 4.1 (Cantor). *(X, τ) is scattered iff $d_\tau^\alpha X = \emptyset$, for some ordinal α .*

Proof. Let $d = d_\tau$. If (X, τ) is scattered then we have $d^\alpha X \supset d^{\alpha+1} X$, for each α such that $d^\alpha X \neq \emptyset$. By cardinality considerations this yields an α such that $d^\alpha X = \emptyset$.

Conversely, suppose $A \subseteq X$ is nonempty. Let α be the least ordinal such that $A \not\subseteq d^\alpha X$. Obviously, α cannot be a limit ordinal, hence $\alpha = \beta + 1$ for some β and there is an $x \in A \setminus d^{\beta+1} X$. Since $A \subseteq d^\beta X$, we also have $x \in d^\beta X$. Since $x \notin d^{\beta+1} X = d(d^\beta X)$, x is isolated in the relative topology of $d^\beta X$, and hence in the relative topology of $A \subseteq d^\beta X$. \dashv

Call the least α such that $d^\alpha X = \emptyset$ the *Cantor–Bendixson rank* of X and denote it by $\rho_\tau(X)$. The *rank function* $\rho_\tau : X \rightarrow \text{On}$ is defined by

$$\rho_\tau(x) := \min\{\alpha : x \notin d_\tau^{\alpha+1}(X)\}.$$

Notice that ρ_τ maps X onto $\rho_\tau(X) = \{\alpha : \alpha < \rho_\tau(X)\}$. Also, $\rho_\tau(x) \geq \alpha$ iff $x \in d_\tau^\alpha X$. We omit the subscript τ whenever there is no danger of confusion.

Example 4.1. Let Ω be an ordinal equipped with its *left topology*. Then $\rho(\alpha) = \alpha$, for all α .

Example 4.2. Let Ω be an ordinal equipped with its *order topology*. Then ρ is the function ℓ defined by

$$\ell(0) = 0; \quad \ell(\alpha) = \beta \text{ if } \alpha = \gamma + \omega^\beta, \text{ for some } \gamma, \beta.$$

By the Cantor normal form theorem, for any $\alpha > 0$, such a β is uniquely defined.

Let (X, τ_X) and (Y, τ_Y) be topological spaces, and let d_X, d_Y denote the corresponding derivative operators. A map $f : X \rightarrow Y$ is called a *d-map* if f is continuous, open and *pointwise discrete*, that is, $f^{-1}(y)$ is a discrete subspace of X for each $y \in Y$. *d*-maps are well-known to satisfy the properties expressed in the following lemma (see [16]).

Lemma 4.2. (i) $f^{-1}(d_Y(A)) = d_X(f^{-1}(A))$, for any $A \subseteq Y$;
(ii) $f^{-1} : (\mathcal{P}(Y), d_Y) \rightarrow (\mathcal{P}(X), d_X)$ is a homomorphism of derivative algebras;
(iii) If f is onto, then $\text{Log}(X, \tau_X) \subseteq \text{Log}(Y, \tau_Y)$.

In fact, (i) is easy to check directly; (ii) follows from (i) and (iii) follows from (ii). Each of the conditions (i) and (ii) is equivalent to f being a *d-map*.

From lemma 4.2(i) we easily obtain the following corollary by transfinite induction.

Corollary 4.3. Suppose $f : X \rightarrow Y$ is a *d-map*. Then, for each ordinal α , $d_X^\alpha X = f^{-1}(d_Y^\alpha Y)$.

The following lemma states that the rank function, when the ordinals are equipped with their left topology, becomes a *d-map*. It is also uniquely characterized by this property.

Lemma 4.4. Let Ω be the ordinal $\rho_\tau(X)$ taken with its left topology. Then

- (i) $\rho_\tau : X \rightarrow \Omega$ is an onto *d-map*;
- (ii) If $f : X \rightarrow \lambda$ is a *d-map*, where λ is an ordinal with its left topology, then $f(X) = \Omega$ and $f = \rho_\tau$.

Proof. Let ρ denote ρ_τ .

- (i) ρ is continuous, because the set $\rho^{-1}[0, \alpha) = X \setminus d^\alpha X$ is open.

ρ being open means that, for each open $U \subseteq X$, whenever $\alpha \in \rho(U)$ and $\beta < \alpha$ one has $\beta \in \rho(U)$. Fix an $x \in U$ such that $\rho(x) = \alpha$. Consider the set $X_\beta := \rho^{-1}(\beta) = d^\beta X \setminus d(d^\beta X)$. For any subset A of a scattered space we have $d(A) = d(A \setminus dA)$, hence $dX_\beta = d(d^\beta X) \subseteq d^\alpha X$. Since $\rho(x) = \alpha$ it follows that $x \in dX_\beta$. Hence $U \cap X_\beta \neq \emptyset$, that is, $\beta \in \rho(U)$.

ρ being pointwise discrete means $X_\alpha = \rho^{-1}(\alpha)$ is discrete, for each α . In fact, $X_\alpha = d^\alpha X \setminus d(d^\alpha X)$ is the set of isolated points of $d^\alpha X$. Thus, it cannot help being discrete.

- (ii) Since f is a *d-map*, by Corollary 4.3 we obtain that $f^{-1}[\alpha, \lambda) = d^\alpha X$, for each $\alpha < \lambda$. Hence, $f^{-1}(\alpha) = \rho^{-1}(\alpha)$, for each $\alpha < \lambda$, that is, $f = \rho$ and $f(X) = \rho(X) = \Omega$. ⊥

Let ρ_X and ρ_Y denote the rank functions of (X, τ_X) and (Y, τ_Y) , respectively.

Corollary 4.5. If $f : X \rightarrow Y$ is a *d-map*, then $\rho_X = \rho_Y \circ f$.

Proof. Clearly, $\rho_Y \circ f : X \rightarrow \Omega$ is a *d-map*. Statement (ii) of the previous lemma yields the result. ⊥

4.2 The d-sum construction

Given a tree T , one can build a new tree by ‘plugging in’ other trees in place of the maximal points (the ‘leaves’) of T . Similarly, given an ordinal α , one can ‘plug in’ new ordinals α_i for each isolated (i.e. non-limit) point $i \in \alpha$ to obtain another ordinal. These constructions of ‘summing up’ spaces or relational structures ‘along’ another space or a relational structure are a special case of a general construction, which proved rather useful in its various manifestations for proving topological completeness theorems. Here we present a variant of this construction, called *d-sum*, which can be used to recursively build both finite irreflexive trees and ordinals. We will point out a particular application of *d-sums* by sketching a proof of Abashidze–Blass theorem. For a more substantial application of the *d-sum* construction, in which the summed spaces are homeomorphic to each other, see [5] (where the corresponding construction is called *d-product*).

Definition 4.6. Let X be a topological space and let $\{Y_j \mid j \in i(X)\}$ be a collection of spaces indexed by the set $i(X)$ of isolated points of X . We uniquely extend it to the collection $\{Y_j \mid j \in X\}$ by letting $Y_j = \{j\}$ for all $j \in d_\tau X$.

We define the *d-sum* (Z, τ_Z) of $\{Y_j\}$ over X (denoted $\sum_{j \in X}^d Y_j$) as follows. The base set is the disjoint union $Z := \bigsqcup_{j \in X} Y_j$. Define the map $\pi : Z \rightarrow X$ by putting $\pi(y) = j$ whenever $y \in Y_j$. Now let the topology τ_Z consist of the sets $V \cup \pi^{-1}(U)$ where V is open in the topological sum $\bigsqcup_{j \in i(X)} Y_j$ and U is open in X . It is not difficult to check that τ_Z qualifies for a topology.

As an immediate application of this construction we present a way of building finite irreflexive trees (called simply *trees* for brevity henceforth) recursively using the *d-sum* construction.

Definition 4.7. (*n-fork*)

Let $\mathfrak{F}_n = (W_n, R_n)$ be defined as follows: $W_n = \{r, w_0, w_1, \dots, w_{n-1}\}$ and $R_n = \{(r, w_i) \mid 0 \leq i < n\}$. We will consider \mathfrak{F}_n equipped with the upset topology w.r.t. R_n .

We view trees, and *n*-forks in particular, as topological spaces.

Definition 4.8. (*trees*)

- (t1) An irreflexive point is a tree.
- (t2) An *n*-fork \mathfrak{F}_n is a tree.
- (t3) The *d-sum* of trees is a tree.

It is straightforward (if somewhat tedious) to check that trees thus obtained are precisely the finite irreflexive trees. Note that the leaves of a tree are the isolated points in the topology. Therefore, taking the *d-sum* of trees T_i over a tree T means simply plugging in T_i ’s in place of the leaves of T . It is worth noting that a tree might be obtained in various different ways as a *d-sum* of other trees.

In a similar fashion the *d-sum* construction can be applied to ordinals to obtain larger ordinals. For example, it is easy to see that summing up ordinals $\omega^n + 1$ along the ordinal $\omega + 1$ produces the ordinal $\omega^{n+1} + 1$. In general, from the well-known theorem characterizing countable ordinals as countable locally compact scattered Hausdorff spaces, it follows that the *d-sum* of countable ordinals is a countable ordinal. In fact, even more general statement is true. Namely, if one

takes a family of ordinals $(\alpha_i)_{i \in \beta}$ such that $\alpha_i = 1$ for limit i , then the ordinal sum $\sum_{i \in \beta} \alpha_i$ is an ordinal that is homeomorphic to the d -sum $\sum_{i \in \beta}^d \alpha_i$. This can be checked directly, by examining the descriptions of neighborhoods in respective topologies. It follows that the d -sum of ordinals is an ordinal.

The last paragraph reveals the intrinsic similarity between the inner structure of trees and ordinals. To fix this similarity, we employ the following general lemma showing that d -sums, in a way, commute with d -maps. More precisely:

Lemma 4.9. *Let X and X' be two spaces and let $\{Y_j \mid j \in i(X)\}$ and $\{Y'_k \mid k \in i(X')\}$ be collections of spaces indexed by $i(X)$ and $i(X')$, respectively. Suppose further that $f : X \rightarrow X'$ is an onto d -map, and for each $j \in i(X)$ there is an onto d -map $f_j : Y_j \rightarrow Y'_{f(j)}$. Then there exists an onto d -map $g : \sum_{j \in X}^d Y_j \rightarrow \sum_{k \in X'}^d Y'_k$.*

Proof. First note that since f is a d -map, $f(j)$ is isolated in X' iff j is isolated in X . Indeed, by openness of f , if $\{j\} \in \tau$, then $\{f(j)\} \in \tau'$. Conversely, if $f(j)$ is isolated, then $f^{-1}f(j)$ is both open and discrete, by continuity and pointwise discreteness of f . Hence, any point in $f^{-1}f(j)$, and j in particular, is isolated in X . For convenience, let us denote $f_* \equiv f|_{d_\tau X}$ and $f^* \equiv f|_{iX}$. It follows that $f^* : i(X) \rightarrow i(X')$ and $f_* : d_\tau X \rightarrow d_{\tau'} X'$ are well-defined onto maps and $f = f^* \cup f_*$. Thus, in particular, the space $Y'_{f(j)}$ in the formulation of the theorem is well-defined.

Take g to be the set-theoretic union $g = f_* \cup \bigcup_{j \in iX} f_j$. We show that g is a d -map. Let π and π' be the ‘projection’ maps associated with $\sum_{j \in X}^d Y_j$ and $\sum_{k \in X'}^d Y'_k$, respectively. To show that g is open, take $W = V \cup \pi^{-1}(U) \in \tau_Z$. Then $g(W) = g(V) \cup g(\pi^{-1}(U))$. That $g(V)$ is open in the topological sum of Y'_k is clear from the openness of the maps f_j . Moreover, from the definition of g and the fact that all f_j are onto it can be easily deduced that $g(\pi^{-1}(U)) = \pi'^{-1}(f(U))$. Since f is an open map, it follows that $g(W)$ is open in τ'_Z . To see that g is continuous, take $W' = V' \cup \pi'^{-1}(U') \in \tau'_Z$. Then $g^{-1}(W') = g^{-1}(U') \cup g^{-1}(\pi'^{-1}(U'))$. Again, the openness of $g^{-1}(U')$ is trivial. It is also easily seen that $g^{-1}(\pi'^{-1}(U')) = \pi^{-1}(f^{-1}(U'))$. It follows that $g^{-1}(W')$ is open in τ_Z . To see that g is pointwise discrete is straightforward, given that f and all the f_j are pointwise discrete. \dashv

In the next section we will use this lemma to prove the Abashidze–Blass theorem.

5 Proof of Abashidze–Blass theorem

The topological completeness theorem proved independently by Abashidze and Blass establishes that each non-theorem of **GL** can be refuted on any ordinal greater than ω^ω . A simplified proof of this result appeared recently in [17, Theorem 3.5]. The crucial part of that proof rests on showing that each tree of depth n is an onto image of an ordinal $\omega^n + 1$ via a d -map [17, Lemma 3.4]. That is probably the simplest and most explicit proof of Abashidze–Blass theorem, however by our approach we aim to illuminate the underlying recursive mechanism, which might offer additional insight. Here is an analog of [17, Lemma 3.4] proved using the d -sum construction:

Lemma 5.1. *For each tree T there exists a countable ordinal $\alpha < \omega^\omega$ and an onto map $f : \alpha \rightarrow T$ such that f is a d -map.*

Proof. [Sketch] The proof proceeds by structural induction. It is clear that the claim is true for a one-point tree. Let us consider now an n -fork \mathfrak{F}_n and build a d -map f from $\omega + 1$ onto \mathfrak{F}_n . Let $f(x) = w_{x \bmod n}$ for $x < \omega$ and let $f(\omega) = r$. It is straightforward to check that f is a d -map. That each tree is an image of an appropriate ordinal via a d -map now follows from lemma 4.9 using induction on the depth of trees. Indeed, suppose the theorem is true for all trees of depth less than n . Take a tree T of depth n . It is clear that T can be presented as a d -sum of trees of strictly smaller depth in various ways. Using the induction hypothesis, each of the smaller trees is an image of a countable ordinal under a d -map. Applying lemma 4.9 and observing that the d -sum of countable ordinals is a countable ordinal produces a countable ordinal α and an onto d -map $f : \alpha \rightarrow T$. By corollary 4.5 the rank of α is equal to the rank of T , that is, to n . It follows that $\alpha < \omega^\omega$, which finishes the proof. \dashv

One of the ways to present the tree T as a d -sum of smaller trees is to ‘prune’ T immediately after the root. Then T is seen as a d -sum of smaller trees along a k -fork for some finite k . This is precisely the approach usually taken in proving the above lemma. However, the d -sum construction allows for greater flexibility. For example, T can be pruned immediately below the leaves, obtaining finitely many forks and a tree of depth $n - 1$. Or indeed, T can be pruned *anywhere*, provided each of the branches of maximal length n is pruned.

And finally, the Abashidze–Blass theorem:

Theorem 5.2 (Abashidze–Blass). *The logic of any ordinal $\alpha \geq \omega^\omega$ is **GL**.*

Proof. Take a non-theorem φ of **GL**. Then φ can be refuted on a finite tree T . By Lemma 5.1 there exists an ordinal $\beta < \omega^\omega$ that maps onto T via a d -map. Since d -maps reflect satisfiability (the contrapositive of Lemma 4.2(iii)), φ can be refuted on β . But β is an open subspace of α . It follows that φ can be refuted on α . \dashv

6 Topological semantics of linearity axioms

For a gödelian theory T consider the subalgebra \mathcal{L}_T^0 of $(\mathcal{L}_T, \Diamond_T)$ generated by \top . If $\text{ch}(\mathcal{L}_T, \Diamond_T) = \infty$, then, clearly, $\text{ch}(\mathcal{L}_T^0, \Diamond_T) = \infty$. In fact, the Magari algebra $(\mathcal{L}_T^0, \Diamond_T)$ is known (see [32]) to have the logic **GL.3** which is obtained from **GL** by adding the following axiom:

$$(.3) \quad \Diamond p \wedge \Diamond q \rightarrow \Diamond(p \wedge q) \vee \Diamond(p \wedge \Diamond q) \vee \Diamond(\Diamond p \wedge q)$$

This is a so called ‘linearity axiom’ and, as the name suggests, its finite rooted Kripke frames are precisely the finite strict linear orders. Since **GL.3** is Kripke complete [23], its topological completeness is almost immediate. However, it is not immediately clear what kind of scattered spaces does the linearity axiom isolate. To characterize GL.3-spaces, let us first simplify the axiom (.3). Let (lin) be the following formula:

$$\Box(\Box^+ p \vee \Box^+ q) \rightarrow \Box p \vee \Box q$$

where $\Box^+ \varphi$ is a shorthand for $\varphi \wedge \Box \varphi$.

Lemma 6.1. *In **GL** the schema (.3) is equivalent to (lin) .*

Proof. To show that $(lin) \vdash_{\mathbf{GL}} (.3)$, witness the following syntactic argument. Observe that the dual form of (lin) looks as follows:

$$\Diamond p \wedge \Diamond q \rightarrow \Diamond(\Diamond^+ p \wedge \Diamond^+ q) \quad (*)$$

where $\Diamond^+ \varphi := \varphi \vee \Diamond \varphi$. Furthermore, an instance of the **GL** axiom looks as follows:

$$\Diamond(\Diamond^+ p \wedge \Diamond^+ q) \rightarrow \Diamond(\Diamond^+ p \wedge \Diamond^+ q \wedge \Box(\Box^+ \neg p \vee \Box^+ \neg q)).$$

By the axiom (lin) we also have:

$$\Box(\Box^+ \neg p \vee \Box^+ \neg q) \rightarrow (\Box \neg p \vee \Box \neg q).$$

So, using the monotonicity of \Diamond we obtain:

$$\Diamond p \wedge \Diamond q \rightarrow \Diamond(\Diamond^+ p \wedge \Diamond^+ q \wedge (\Box \neg p \vee \Box \neg q)).$$

By Boolean logic we clearly have:

$$\Diamond^+ p \wedge \Diamond^+ q \leftrightarrow (p \wedge q) \vee (p \wedge \Diamond q) \vee (\Diamond p \wedge q) \vee (\Diamond p \wedge \Diamond q) \quad (**)$$

and

$$(\Box \neg p \vee \Box \neg q) \leftrightarrow \neg(\Diamond p \wedge \Diamond q).$$

Using these, together with the monotonicity of \Diamond we finally arrive at:

$$\Diamond p \wedge \Diamond q \rightarrow \Diamond((p \wedge q) \vee (p \wedge \Diamond q) \vee (\Diamond p \wedge q)),$$

which is equivalent to $(.3)$ since \Diamond distributes over \vee .

To show the converse, we observe that $(.3)$ implies (lin) even in the system **K**. Indeed, the formula $(*)$, which is the dual form of (lin) , can be rewritten, using $(**)$ and the distribution of \Diamond over \vee as follows:

$$\Diamond p \wedge \Diamond q \rightarrow \Diamond(p \wedge q) \vee \Diamond(p \wedge \Diamond q) \vee \Diamond(\Diamond p \wedge q) \vee \Diamond(\Diamond p \wedge \Diamond q),$$

which is clearly a weakening of $(.3)$. Therefore $(.3) \vdash_{\mathbf{GL}} (lin)$. \dashv

It follows that a scattered space is a GL.3-space iff it validates (lin) . To characterize such spaces, consider the following definition.

Definition 6.2. Call a scattered space *primal* if for each $x \in X$ and $U, V \in \tau$, $\{x\} \cup U \cup V \in \tau$ implies $\{x\} \cup U \in \tau$ or $\{x\} \cup V \in \tau$.

In other words, X is primal if the collection of punctured neighborhoods of each non-isolated point is a prime filter in the Heyting algebra τ .

Theorem 6.3. *Let X be a scattered space. Then $X \models (lin)$ iff X is primal.*

Proof. Let X be a scattered space together with a valuation v . Let $P := v(p)$ and $Q := v(q)$ denote the truth-sets of p and q respectively. Then the truth sets of $\Box^+ p$ and $\Box^+ q$ are $I_\tau P$ and $I_\tau Q$, where I_τ is the interior operator of X . We write $x \models \varphi$ for $X, x \models_v \varphi$ (as introduced in section 3).

Suppose X is primal and for some valuation $x \models \Box(\Box^+ p \vee \Box^+ q)$. Then there exists an open neighborhood W of x such that $W \setminus \{x\} \models \Box^+ p \vee \Box^+ q$. In other words, $W \setminus \{x\} \subseteq I_\tau P \cup I_\tau Q$. Let $U = W \cap I_\tau P \in \tau$ and $V = W \cap I_\tau Q \in \tau$. Then

$\{x\} \cup U \cup V = W \in \tau$. It follows that either $\{x\} \cup U \in \tau$ or $\{x\} \cup V \in \tau$. Hence $x \models \Box p$ or $x \models \Box q$. This proves that $X \models (lin)$.

Suppose now X is not primal. Then there exist $x \in X$ and $U, V \in \tau$ such that $\{x\} \cup U \cup V \in \tau$, but $\{x\} \cup U \notin \tau$ and $\{x\} \cup V \notin \tau$. Take a valuation such that $P = U$ and $Q = V$. Then clearly $x \models \Box(\Box^+ p \vee \Box^+ q)$. However, neither $x \models \Box p$ nor $x \models \Box q$ is true. Indeed, if, for example, $x \models \Box p$, then there exists an open neighborhood W of x such that $W \setminus \{x\} \subseteq P = U$. But then $\{x\} \cup U = W \cup U \in \tau$, which is a contradiction. This shows that $X \not\models (lin)$. \dashv

The primal scattered spaces are very close to *maximal scattered* spaces of [5]. In fact, each maximal scattered space is primal, but there are primal spaces which are not maximal. It follows that the logic of maximal scattered spaces is **GL.3**.

7 GLP-algebras and polymodal provability logic

A natural generalization of provability logic **GL** to a language with infinitely many modalities $\langle 0 \rangle, \langle 1 \rangle, \dots$ has been introduced as early as in 1986 by Giorgi Japaridze [35]. He interpreted $\langle 1 \rangle \varphi$ as an arithmetical statement expressing the ω -consistency of φ over a given gödelian theory T .² Similarly, $\langle 2 \rangle \varphi$ was interpreted as the consistency of the extension of $T + \varphi$ by two nested applications of ω -rule, and so on.

While the logic of each of the individual modalities $\langle n \rangle$ over Peano arithmetic was known to coincide with **GL** by a relatively straightforward extension of the Solovay Theorem [19], Japaridze found a complete axiomatization of the *joint* logic of the modalities $\langle n \rangle$, for all $n \in \omega$. This result involved considerable technical difficulties and lead to one of the first genuine extensions of Solovay's arithmetical fixed-point construction. Later, Japaridze's work has been simplified and extended by Konstantin Ignatiev [34] and George Boolos [20]. In particular, Ignatiev showed that **GLP** is complete for more general sequences of 'strong' provability predicates in arithmetic and analyzed the variable-free fragment of **GLP**. Boolos included a treatment of **GLP** (for the somewhat simpler case of just two modalities) in his popular book on provability logic [21].

More recently, **GLP** has found interesting applications in proof-theoretic analysis of arithmetic [6, 7, 2, 9] which stimulated some further interest in the study of modal-logical properties of **GLP** [22, 15, 33, 11]. For such applications, the algebraic language appears to be more natural and a different choice of the interpretation of the provability predicates is needed. The relevant structures have been introduced in [6] under the name *graded provability algebras*.

Recall that an arithmetical formula is called Π_n if it can be obtained from a formula containing only bounded quantifiers $\forall x \leq t$ and $\exists x \leq t$ by a prefix of n alternating blocks of quantifiers starting from \forall . Arithmetical Σ_n -formulas are defined dually.

Let T be a gödelian theory. T is called *n-consistent* if T together with all true arithmetical Π_n -sentences is consistent. (Alternatively, T is *n-consistent* iff every Σ_n -sentence provable in T is true.) Let $n\text{-Con}(T)$ denote a natural arithmetical

²A gödelian theory U is ω -consistent if its extension by unnested applications of the ω -rule $U' := U + \{\forall x \varphi(x) : \forall n U \vdash \varphi(\underline{n})\}$ is consistent.

formula expressing the n -consistency of T (it can be defined using the standard Π_n -definition of truth for Π_n -sentences in arithmetic). Since we assume T to be recursively enumerable, it is easy to check that the formula $n\text{-Con}(T)$ itself belongs to the class Π_{n+1} .

The n -consistency formula induces an operator $\langle n \rangle_T$ acting on the Lindenbaum–Tarski algebra \mathcal{L}_T :

$$\langle n \rangle_T : [\varphi] \longmapsto [n\text{-Con}(T + \varphi)].$$

The dual n -provability operators are defined by $[n]_T x = \neg \langle n \rangle_T x$, for all $x \in \mathcal{L}_T$. Since every true Π_n -sentence is assumed to be an axiom for n -provability, we notice that every true Σ_{n+1} -sentence must be n -provable. Moreover, this latter fact is formalizable in T , so we obtain the following lemma. (By the abuse of notation we denote by $[n]_T \varphi$ also the arithmetical formula expressing the n -provability of φ in T .)

Lemma 7.1. *For each true Σ_{n+1} -formula $\sigma(x)$,*

$$T \vdash \forall x (\sigma(x) \rightarrow [n]_T \sigma(\underline{x})).$$

As a corollary we obtain the following basic observation probably due to Smoryński (see [45]).

Proposition 7.2. *For each $n \in \omega$, the structure $(\mathcal{L}_T, \langle n \rangle_T)$ is a Magari algebra.*

A proof of this fact consists of verifying the Bernays–Löb derivability conditions for $[n]_T$ in T and of deducing from them, in the usual way, an analog of Löb’s theorem for $[n]_T$.

The structure $(\mathcal{L}_T, \{\langle n \rangle_T : n \in \omega\})$ is called the *graded provability algebra of T* or the *GLP-algebra of T* . Apart from the identities inherited from the structure of Magari algebras for each $\langle n \rangle$ it satisfies the following principles, for all $m < n$:

P1. $\langle m \rangle x \leq [n] \langle m \rangle x$;

P2. $\langle n \rangle x \leq \langle m \rangle x$.

The validity of P1 follows from Lemma 7.1, because the formula $\langle m \rangle_T \varphi$, for any φ , belongs to the class Π_{m+1} . P2 holds, since $\langle n \rangle_T \varphi$ asserts the consistency of a stronger theory than $\langle m \rangle_T \varphi$, for $m < n$.

In general, we call a *GLP-algebra* a structure $(M, \{\langle n \rangle : n \in \omega\})$ such that each $(M, \langle n \rangle)$ is a Magari algebra and conditions P1, P2 (that are equivalent to identities) are satisfied for all $x \in M$.

At this point it is worth noticing that P1 has an equivalent form (modulo the other identities) that has proved to be quite useful for various applications of GLP-algebras.

Lemma 7.3. *Modulo the identities of Magari algebras and P2, condition P1 is equivalent to*

P1'. $\langle n \rangle y \wedge \langle m \rangle x = \langle n \rangle (y \wedge \langle m \rangle x)$, for all $m < n$.

Proof. First, we prove P1'. We have $y \wedge \langle m \rangle x \leq y$, hence

$$\langle n \rangle (y \wedge \langle m \rangle x) \leq \langle n \rangle y.$$

Similarly, by P2 and transitivity,

$$\langle n \rangle(y \wedge \langle m \rangle x) \leq \langle n \rangle \langle m \rangle x \leq \langle m \rangle \langle m \rangle x \leq \langle m \rangle x.$$

Hence, $\langle n \rangle(y \wedge \langle m \rangle x) \leq \langle n \rangle y \wedge \langle m \rangle x$. In the other direction, by P1,

$$\langle n \rangle y \wedge \langle m \rangle x \leq \langle n \rangle y \wedge [n] \langle m \rangle x.$$

However, as in any modal algebra, we also have $\langle n \rangle y \wedge [n] z \leq \langle n \rangle(y \wedge z)$. It follows that

$$\langle n \rangle y \wedge [n] \langle m \rangle x \leq \langle n \rangle(y \wedge \langle m \rangle x).$$

Thus, P1' is proved.

To infer P1 from P1' it is sufficient to prove that

$$\langle m \rangle x \wedge \neg[n] \langle m \rangle x = \perp.$$

We have that $\neg[n] \langle m \rangle x = \langle n \rangle \neg \langle m \rangle x$. Therefore, by P1',

$$\langle m \rangle x \wedge \langle n \rangle \neg \langle m \rangle x = \langle n \rangle(\neg \langle m \rangle x \wedge \langle m \rangle x) = \langle n \rangle \perp = \perp,$$

as required. ⊥

An equivalent formulation of Japaridze's arithmetical completeness theorem is that any identity of $(\mathcal{L}_T, \{\langle n \rangle_T : n \in \omega\})$ follows from the identities of GLP-algebras [35]. It is somewhat strengthened to the current formulation in [34, 13].

Theorem 7.4 (Japaridze). *Suppose T is gödelian, T contains Peano arithmetic, and $\text{ch}(\mathcal{L}_T, \langle n \rangle_T) = \infty$, for each $n < \omega$. Then, an identity holds in $(\mathcal{L}_T, \{\langle n \rangle_T : n \in \omega\})$ iff it holds in all GLP-algebras.*

We note that the condition $\text{ch}(\mathcal{L}_T, \langle n \rangle_T) = \infty$, for each $n \in \omega$, is equivalent to $T + n\text{-Con}(T)$ being consistent, for each $n \in \omega$, and is clearly necessary for the validity of Japaridze's theorem.

The logic of all GLP-algebras can also be axiomatized as a Hilbert-style calculus. The corresponding system **GLP** was originally introduced by Japaridze. **GLP** is formulated in the language of propositional logic enriched by modalities $[n]$, for all $n \in \omega$. The axioms of **GLP** are those of **GL**, formulated for each $[n]$, as well as the two analogs of P1 and P2, for all $m < n$:

$$\text{P1. } \langle m \rangle \varphi \rightarrow [n] \langle m \rangle \varphi;$$

$$\text{P2. } [m] \varphi \rightarrow [n] \varphi.$$

The inference rules of **GLP** are modus ponens and $\varphi/[n] \varphi$, for each $n \in \omega$.

For any modal formula φ , **GLP** $\vdash \varphi$ iff the identity $t_\varphi = \top$ holds in all **GLP**-algebras. Hence, **GLP** coincides with logic of all GLP-algebras as well as with the logic of the GLP-algebra of T , for any theory T such that $T + n\text{-Con}(T)$ is consistent, for each $n < \omega$.

8 GLP-spaces

Topological semantics for **GLP** has been first considered in [14]. The main difficulty in the modal-logical study of **GLP** comes from the fact that it is incomplete with respect to its relational semantics; that is, **GLP** is the logic of no class of *frames*. Even though a suitable class of relational *models* for which **GLP** is sound and complete was developed in [11], these models are sufficiently complicated and not so easy to handle. So, it is natural to consider the generalization of the topological semantics we have for **GL**. As it turns out, topological semantics provides another natural class of GLP-algebras which is interesting in its own right, as well as because of its analogy with the proof-theoretic GLP-algebras.

As before, we are interested in GLP-algebras of the form $(\mathcal{P}(X), \{\langle n \rangle : n \in \omega\})$ where $\mathcal{P}(X)$ is the boolean algebra of subsets of a given set X . Since each $(\mathcal{P}(X), \langle n \rangle)$ is a Magari algebra, the operator $\langle n \rangle$ is the derivative operator with respect to some uniquely defined scattered topology on X . Thus, we come to the following definition [14].

A polytopological space $(X, \{\tau_n : n \in \omega\})$ is called a *GLP-space* if the following conditions hold, for each $n \in \omega$:

- D0. (X, τ_n) is a scattered topological space;
- D1. For each $A \subseteq X$, $d_{\tau_n}(A)$ is τ_{n+1} -open;
- D2. $\tau_n \subseteq \tau_{n+1}$.

We notice that the last two conditions directly correspond to conditions P1 and P2 of GLP-algebras. By a *GLP_m-space* we mean a space $(X, \{\tau_n : n < m\})$ satisfying conditions D0–D2 for the first m topologies.

Proposition 8.1. (i) *If $(X, \{\tau_n : n \in \omega\})$ is a GLP-space then the structure $(\mathcal{P}(X), \{d_{\tau_n} : n \in \omega\})$ is a GLP-algebra.*

(ii) *If $(\mathcal{P}(X), \{\langle n \rangle : n \in \omega\})$ is a GLP-algebra, then there are uniquely defined topologies $\{\tau_n : n \in \omega\}$ on X such that $(X, \{\tau_n : n \in \omega\})$ is a GLP-space and $\langle n \rangle = d_{\tau_n}$, for each $n < \omega$.*

Proof. (i) Suppose $(X, \{\tau_n : n \in \omega\})$ is a GLP-space. Let $d_n := d_{\tau_n}$ denote the corresponding derivative operators and let \tilde{d}_n denote its dual $\tilde{d}_n(A) := X \setminus d_n(X \setminus A)$. By Theorem 2.1 $(\mathcal{P}(X), d_n)$ is a Magari algebra, for each $n \in \omega$. Notice that $A \in \tau_n$ iff $A \subseteq \tilde{d}_n A$. If $m < n$ then $d_m A \in \tau_n$, hence $d_m A \subseteq \tilde{d}_n d_m A$, hence P1 holds. Since $\tau_n \subseteq \tau_{n+1}$ we have $d_{n+1} A \subseteq d_n A$ hence P2 holds.

(ii) Let $(\mathcal{P}(X), \{\langle n \rangle : n \in \omega\})$ be a GLP-algebra. Since each of the algebras $(\mathcal{P}(X), \langle n \rangle)$ is Magari, by Theorem 2.4 a scattered topology τ_n on X is defined for which $\langle n \rangle = d_{\tau_n}$. In fact, we have $U \in \tau_n$ iff $U \subseteq [n]U$. We check that conditions D1 and D2 are met.

Suppose A is τ_n -closed, that is, $\langle n \rangle A \subseteq A$. Then $\langle n+1 \rangle A \subseteq \langle n \rangle A \subseteq A$ by P2. Hence, U is τ_{n+1} -closed. Thus, $\tau_n \subseteq \tau_{n+1}$.

By P1, for any set A we have $\langle n \rangle A \subseteq [n+1]\langle n \rangle A$. Hence, $d_{\tau_n}(A) = \langle n \rangle A \in \tau_{n+1}$. Thus, $(X, \{\tau_n : n \in \omega\})$ is a GLP-space. \dashv

9 Derivative topology and generated GLP-spaces

To obtain examples of GLP-spaces let us first consider the case of two modalities. The following basic example is due to Esakia (private communication).

Example 9.1. Consider a bitopological space $(\Omega; \tau_0, \tau_1)$ where Ω is an ordinal, τ_0 is its left topology, and τ_1 is its order topology. Esakia noticed that this space is a model of the bimodal fragment of **GLP**, that is, in our terminology, a GLP_2 -space. In fact, for any $A \subseteq \Omega$ the set $d_0(A) = (\min A, \Omega)$ is an open interval, whenever A is not empty. Hence, condition D1 holds (the other two conditions are immediate). Esakia also noticed that such spaces can never be complete for **GLP**, as the linearity axiom (.3) holds for $\langle 0 \rangle$.

In general, to define GLP_n -spaces for $n > 1$, we introduce an operation $\tau \mapsto \tau^+$ on topologies on a given set X . This operation plays a central role in the study of GLP-spaces.

Given a topological space (X, τ) , let τ^+ be the coarsest topology containing τ such that each set of the form $d_\tau(A)$, with $A \subseteq X$, is open in τ^+ . Thus, τ^+ is generated by τ and $\{d_\tau(A) : A \subseteq X\}$. Clearly, τ^+ is the coarsest topology on X such that $(X; \tau, \tau^+)$ is a GLP_2 -space. Sometimes we call τ^+ the *derivative topology* of (X, τ) .

Getting back to Esakia's example, it is easy to verify that, on any ordinal Ω , the derivative topology of the left topology coincides with the order topology. (In fact, any open interval is an intersection of a downset and an open upset.)

Example 9.2. Even though we are mainly interested in scattered spaces, the derivative topology makes sense for arbitrary spaces. The reader can check that if τ is the *coarsest* topology on a set X (whose open sets are just X and \emptyset), then τ^+ is the *cofinite* topology on X (whose open sets are exactly the cofinite subsets of X). On the other hand, if τ is the cofinite topology then $\tau^+ = \tau$. We note that the logic of the cofinite topology on an infinite set is **KD45** (see [47]).

For scattered spaces, τ^+ is always strictly finer than τ , unless τ is discrete. We present a proof using the language of Magari algebras.

Proposition 9.1. *Suppose (X, τ) is scattered. Then $d_\tau(X)$ is not open, unless $d_\tau(X) = \emptyset$.*

Proof. The set $d_\tau(X)$ corresponds to the element $\Diamond \top$ in the associated Magari algebra; $d_\tau(X)$ being open means $\Diamond \top \leq \Box \Diamond \top$. By Löb's principle we have $\Box \Diamond \top \leq \Box \perp = \neg \Diamond \top$. Hence, $\Diamond \top \leq \neg \Diamond \top$, that is, $\Diamond \top = \perp$. This means $d_\tau(X) = \emptyset$, as required. \dashv

In general, we will see later that τ^+ can be much finer than τ . Notice that if τ is T_d then each set of the form $d_\tau(A)$ is τ -closed. Hence, it will be clopen in τ^+ . Thus, τ^+ is obtained by adding to τ new clopen sets. In particular, τ^+ will be zero-dimensional if so is τ .³

Iterating the plus operation delivers us a GLP-space. Let (X, τ) be a scattered space. Define: $\tau_0 := \tau$ and $\tau_{n+1} := \tau_n^+$. Then $(X, \{\tau_n : n \in \omega\})$ is a GLP-space that will be called the GLP-space *generated from* (X, τ) , or simply the *generated GLP-space*.

³Recall that a topological space is zero-dimensional if it has a base of clopen sets.

Thus, from any scattered space we can always produce a GLP-space in a natural way. The question is whether this space will be nontrivial, that is, whether we can guarantee that the topologies τ_n are non-discrete.

In fact, the next observation shows that for many natural τ already the topology τ^+ will be discrete. Recall that a topological space X is *first-countable* if every point $x \in X$ has a countable basis of open neighborhoods.

Proposition 9.2. *If (X, τ) is Hausdorff and first-countable, then τ^+ is discrete.*

Proof. It is easy to see that if (X, τ) is first-countable and Hausdorff, then every point $a \in d_\tau(X)$ is a (unique) limit point of a countable sequence of points $A = \{a_n\}_{n \in \omega}$. Hence, there is a set $A \subseteq X$ such that $d_\tau(A) = \{a\}$. By D1 this means that $\{a\}$ is τ^+ -open. \dashv

Thus, if τ is the order topology on a countable ordinal, then τ^+ is discrete. The same holds, for example, if τ is the (non-scattered) topology of the real line.

We remark that the left topology τ on any countable ordinal $> \omega$ yields an example of a first-countable space such that τ^+ is non-discrete. In the following section we will also see that, if τ is the order topology on any ordinal $> \omega_1$, then τ^+ is non-discrete (ω_1 is its least non-isolated point). However, we do not have any topological characterization of spaces (X, τ) such that τ^+ is discrete. (See, however, Proposition 10.3, which provides a characterization in terms of d -reflection.)

It is a natural question to ask what kind of separation properties is τ^+ guaranteed to have, for an arbitrary (scattered) topology τ . In fact, for τ^+ we can infer a bit more separation than for an arbitrary scattered topology. Recall that a topological space X is T_1 if, for any two different points $a, b \in X$, there is an open set U such that $a \in U$ and $b \notin U$.

Proposition 9.3. *Let (X, τ) be any topological space. Then (X, τ^+) is T_1 .*

Proof. Let $a, b \in X$, $a \neq b$. We must show that there is an open set A such that $a \in A$ and $b \notin A$. Consider the set $B := d_\tau(\{b\})$, which is open in τ^+ . If $a \in B$ then we are done, because $b \notin d_\tau(\{b\}) = B$. Otherwise, if $a \notin B$, neither does a belong to the closure of $\{b\}$ (which is simply $\{b\} \cup d_\tau(\{b\})$). It follows that the complement of $\{b\} \cup B$ is the required open set. \dashv

The following example shows that, in general, τ^+ need not always be Hausdorff.

Example 9.3. Let (X, \prec) be a strict partial ordering on $X := \omega \cup \{a, b\}$ where ω is taken with its natural order, a and b are \prec -incomparable, and $n \prec a, b$, for all $n \in \omega$. Let τ be the left topology on (X, \prec) . Since \prec is well-founded, τ is scattered.

Notice that $d_\tau(A) = \{x \in X : \exists y \in A \ y \prec x\}$. Hence, if A intersects ω , then $d_\tau(A)$ contains an end-segment of ω . Otherwise, $d_\tau(A) = \emptyset$. It follows that a base of open neighborhoods of a in τ^+ consists of sets of the form $I \cup \{a\}$ where I is an end-segment of ω . Similarly, sets of the form $I \cup \{b\}$ are a base of open neighborhoods of b . But any two such sets have a non-empty intersection.

10 d -reflection

In the next section we are going to describe in some detail the GLP-space generated from the left topology on the ordinals. Strikingly, we will see that it naturally leads to some of the central notions of combinatorial set theory, such as Mahlo operation and stationary reflection. In fact, part of our analysis can be easily stated using the language of modal logic for arbitrary generated GLP-spaces. In this section we provide a necessary setup and characterize the topologies of a generated GLP-space in terms of what we call *d-reflection*.⁴

Throughout this section we will assume that (X, τ) is a T_d space and let $d = d_\tau$. Thus, $(\mathcal{P}(X), d)$ is a K4-algebra.

Definition 10.1. A point $a \in X$ is called *d-reflexive* if $a \in dX$ and, for each $A \subseteq X$,

$$a \in dA \Rightarrow a \in d(dA).$$

In modal logic terms this means that the following formula is valid at $a \in X$, for any evaluation of the variable p :

$$X, a \models \Diamond \top \wedge (\Diamond p \rightarrow \Diamond \Diamond p).$$

Similarly, a point $a \in X$ is called *m-fold d-reflexive* if $a \in dX$ and, for each $A_1, \dots, A_m \subseteq X$,

$$a \in dA_1 \cap \dots \cap dA_m \Rightarrow a \in d(dA_1 \cap \dots \cap dA_m).$$

2-fold *d-reflexive* points will also be called *doubly d-reflexive* points. Modal logically, $a \in X$ is doubly *d-reflexive* iff

$$X, a \models \Diamond \top \wedge (\Diamond p \wedge \Diamond q \rightarrow \Diamond(\Diamond p \wedge \Diamond q)).$$

Lemma 10.2. *Each doubly d-reflexive point $x \in X$ is m-fold d-reflexive, for any finite m .*

Proof. The argument goes by induction on $m \geq 2$. Suppose $x \in dA_1 \cap \dots \cap dA_{m+1}$, then $x \in dA_1 \cap \dots \cap dA_m$ and $x \in dA_{m+1}$. By induction hypothesis,

$$x \in d(dA_1 \cap \dots \cap dA_m)$$

and by 2-fold reflection

$$x \in d(d(dA_1 \cap \dots \cap dA_m) \cap dA_{m+1}).$$

However, by T_d property

$$d(dA_1 \cap \dots \cap dA_m) \subseteq dA_1 \cap \dots \cap dA_m,$$

hence

$$x \in d(dA_1 \cap \dots \cap dA_m \cap dA_{m+1}),$$

as required. ⊢

⁴Curiously, the reader may notice that the notion of *reflection principle* as used in provability logic and formal arithmetic matches very nicely the notions such as *stationary reflection* in set theory. (As far as we know, the two terms have evolved completely independently from one another.)

Proposition 10.3. *Let (X, τ) be a T_d -space. A point $x \in X$ is doubly d -reflexive iff x is a limit point of (X, τ^+) .*

Proof. For the (if) direction, we give an argument in the algebraic format. In fact, it is sufficient to show the following inequality in the algebra of (X, τ) , for any elements $p, q \subseteq X$:

$$\langle 1 \rangle \top \wedge \langle 0 \rangle p \wedge \langle 0 \rangle q \leq \langle 0 \rangle (\langle 0 \rangle p \wedge \langle 0 \rangle q).$$

Notice that by Proposition 7.3,

$$\langle 1 \rangle \top \wedge \langle 0 \rangle p = \langle 1 \rangle (\top \wedge \langle 0 \rangle p) = \langle 1 \rangle \langle 0 \rangle p.$$

Hence, we obtain using P1' once again:

$$\langle 1 \rangle \top \wedge \langle 0 \rangle p \wedge \langle 0 \rangle q = \langle 1 \rangle \langle 0 \rangle p \wedge \langle 0 \rangle q = \langle 1 \rangle (\langle 0 \rangle p \wedge \langle 0 \rangle q).$$

The latter formula can be weakened to $\langle 0 \rangle (\langle 0 \rangle p \wedge \langle 0 \rangle q)$ by P2, as required.

For the (only if) direction, it is sufficient to show that each doubly d -reflexive point of (X, τ) is a limit point of τ^+ . Suppose x is doubly d -reflexive. By Lemma 10.2, x is m -fold d -reflexive.

Any basic open subset of τ^+ has the form

$$U := A_0 \cap dA_1 \cap \cdots \cap dA_m,$$

where $A_0 \in \tau$. Assume $x \in U$, we have to find a point $y \neq x$ such that $y \in U$.

Since $x \in dA_1 \cap \cdots \cap dA_m$, by m -fold d -reflexivity we obtain $x \in d(dA_1 \cap \cdots \cap dA_m)$. Since A_0 is an open neighborhood of x , there is a $y \in A_0$ such that $y \neq x$ and $y \in dA_1 \cap \cdots \cap dA_m$. Hence $y \in U$ and $y \neq x$, as required. \dashv

Let d^+ denote the derivative operator associated with τ^+ . We obtain the following characterization of derived topology in terms of neighborhoods.

Proposition 10.4. *A subset $U \subseteq X$ contains a τ^+ -neighborhood of $x \in X$ iff one of the following two cases holds:*

- (i) x is not doubly d -reflexive and $x \in U$;
- (ii) x is doubly d -reflexive and there is an $A \in \tau$ and a B such that $x \in A \cap dB \subseteq U$.

Proof. Since (i) ensures that x is τ^+ -isolated, each condition is clearly sufficient for U to contain a τ^+ -neighborhood of x . To prove the converse, assume that U contains a τ^+ -neighborhood of x . This means

$$x \in A \cap dA_1 \cap \cdots \cap dA_m \subseteq U,$$

for some A, A_1, \dots, A_m with $A \in \tau$.

If x is τ^+ -isolated, condition (i) holds. Otherwise, $x \in d^+X$. Let $B := dA_1 \cap \cdots \cap dA_m$. Since B is closed in τ we have $dB \subseteq B$, hence $A \cap dB \subseteq U$. It remains us to show that $x \in A \cap dB$. By Lemma 7.3, $B \cap d^+X = d^+B \subseteq dB$. Hence, $x \in A \cap B \cap d^+X \subseteq A \cap dB$. \dashv

Remark 10.1. Since in (ii) of the previous lemma A is open, we have $A \cap dB = A \cap d(A \cap B)$, for any B . Hence, we can assume $B \subseteq A$.

Corollary 10.5. *Let (X, τ) be a T_d -space. Then, for all $x \in X$ and $A \subseteq X$, $x \in d^+A$ iff the following two conditions hold:*

- (i) *x is doubly d -reflexive;*
- (ii) *For all $B \subseteq X$, $x \in dB \Rightarrow x \in d(A \cap dB)$.*

The second condition is similar to x being d -reflexive, however dB is now required to reflect from x to a point of A .

Proof. The fact that (i) and (ii) are necessary is proved using Proposition 10.3 and the inequality

$$d^+A \cap dB = d^+(A \cap dB) \subseteq d(A \cap dB).$$

We prove that (i) and (ii) are sufficient. Assume $x \in U \in \tau^+$. By Proposition 10.4 we can assume that U has the form $V \cap dB$ where $V \in \tau$. By (ii), from $x \in dB$ we obtain $x \in d(A \cap dB)$. Hence, there is a $y \neq x$ such that $y \in V$ and $y \in A \cap dB$. It follows that $y \in A$ and $y \in V \cap dB = U$. \dashv

11 The ordinal GLP-space

Here we discuss the GLP-space generated from the left topology on the ordinals, that is, the GLP-space $(\Omega; \{\tau_n : n \in \omega\})$ with τ_0 the left topology on Ω and $\tau_{n+1} = \tau_n^+$, for each $n \in \omega$. It is convenient to think of Ω as the class of all ordinals, even though some readers might feel safer with Ω being a large fixed ordinal. Our basic findings are summarized in the following table, to which we provide extended comments below.

The rows of the table correspond to topologies τ_n . The first column contains the name of the topology (the first two are standard, the third one is introduced in [14], the fourth one is introduced here). The second column indicates the first limit point of τ_n , which is denoted θ_n . The last column describes the derivative operator associated with τ_n . We note that θ_3 is a large cardinal which is sometimes referred to as *the first cardinal reflecting for pairs of stationary sets* (see below), but we know no special notation for this cardinal.

	name	θ_n	$d_n(A)$
τ_0	left	1	$\{\alpha : A \cap \alpha \neq \emptyset\}$
τ_1	order	ω	$\{\alpha \in \text{Lim} : A \cap \alpha \text{ is unbounded in } \alpha\}$
τ_2	club	ω_1	$\{\alpha : \text{cf}(\alpha) > \omega \text{ and } A \cap \alpha \text{ is stationary in } \alpha\}$
τ_3	Mahlo	θ_3	$\dots \dots$

We have already seen that the derivative topology of the left topology is exactly the order topology. Therefore, basic facts related to the first two rows of the table are rather clear. We turn to the next topology τ_2 .

11.1 Club topology

To characterize τ_2 we apply Proposition 10.4, hence it is useful to see what corresponds to the notion of doubly d -reflexive point of the interval topology.

Recall that the *cofinality* $\text{cf}(\alpha)$ of a limit ordinal α is the least order type of a cofinal subset of α ; $\text{cf}(\alpha) := 0$ if $\alpha \notin \text{Lim}$. An ordinal α is *regular* if $\text{cf}(\alpha) = \alpha$.

Lemma 11.1. *For any ordinal α , α is d_1 -reflexive iff α is doubly d_1 -reflexive iff $\text{cf}(\alpha) > \omega$.*

Proof. d_1 -reflexivity of α implies $\text{cf}(\alpha) > \omega$. In fact, d_1 -reflexivity of α means that $\alpha \in \text{Lim}$ and, for all $A \subseteq \alpha$, if A is cofinal in α , then $d_1(A)$ is cofinal in α . If $\text{cf}(\alpha) = \omega$ then there is an increasing sequence $(\alpha_n)_{n \in \omega}$ such that $\sup\{\alpha_n : n \in \omega\} = \alpha$. Then, for $A := \{\alpha_n : n \in \omega\}$ we obviously have $d_1(A) = \{\alpha\}$, hence A violates the reflexivity property.

Now we show that $\text{cf}(\alpha) > \omega$ implies α is doubly d_1 -reflexive. Suppose $\text{cf}(\alpha) > \omega$ and $A, B \subseteq \alpha$ are both cofinal in α . We show that $d_1 A \cap d_1 B$ is cofinal in α . Assume $\beta < \alpha$. Using the cofinality of A, B we can construct an increasing sequence $(\gamma_n)_{n \in \omega}$ above β such that $\gamma_n \in A$, for even n , and $\gamma_n \in B$ for odd n . Let $\gamma := \sup\{\gamma_n : n < \omega\}$. Obviously, both A and B are cofinal in γ whence $\gamma \in d_1 A \cap d_1 B$. Since $\text{cf}(\alpha) > \omega$ and $\text{cf}(\gamma) = \omega$, we have $\gamma < \alpha$. \dashv

Corollary 11.2. *Limit points of τ_2 are exactly the ordinals of uncountable cofinality.*

It turns out that topology τ_2 is strongly related to the well-known concept of *club filter* in set theory.

Let α be a limit ordinal. A subset $C \subseteq \alpha$ is called a *club* in α if C is closed in the order topology of α and unbounded in α . The filter on α generated by all clubs in α is called the *club filter*.

Proposition 11.3. *Assume $\text{cf}(\alpha) > \omega$. The following statements are equivalent:*

- (i) U contains a τ_2 -neighborhood of α ;
- (ii) There is a $B \subseteq \alpha$ such that $\alpha \in d_1 B \subseteq U$;
- (iii) $\alpha \in U$ and U contains a club in α ;
- (iv) $\alpha \in U$ and $U \cap \alpha$ belongs to the club filter on α .

Proof. Statement (ii) implies (iii), since $\alpha \cap d_1 B$ is a club in α , whenever $\alpha \in d_1 B$. Statement (iii) implies (iv), for obvious reasons.

Statement (iv) implies (i). If C is a club in α , then $C \cup \{\alpha\}$ contains a τ_2 -neighborhood $d_1 C$ of α . Indeed, $d_1 C$ is τ_2 -open, contains α , and $d_1 C \subseteq C \cup \{\alpha\}$ since C is τ_1 -closed in α .

Statement (i) implies (ii). Assume U contains a τ_2 -neighborhood of α . Since $\text{cf}(\alpha) > \omega$, by Lemma 11.1 and Proposition 10.4 there is an $A \in \tau_1$ and a B_1 such that $\alpha \in A \cap d_1 B_1 \subseteq U$. Since A is a τ_1 -neighborhood of α , by Proposition 10.4 there are $A_0 \in \tau_0$ and a B_0 such that $\alpha \in A_0 \cap d_0 B_0$. Since τ_0 is the left topology, we can assume that A_0 is the minimal τ_0 -neighborhood $[0, \alpha]$ of α . Besides, we have $\alpha \in d_0 B_0 \cap d_1 B_1 = d_1(B_1 \cap d_0 B_0) \subseteq U$. Since $[0, \alpha]$ is τ_1 -clopen, $d_1(C \cap \alpha) = [0, \alpha] \cap d_1 C$ for any C , so we can take $B_1 \cap d_0 B_0 \cap \alpha$ for B . \dashv

Definition 11.4. The *club topology* on Ω is the unique topology such that

- If $\text{cf}(\alpha) \leq \omega$ then α is an isolated point;
- If $\text{cf}(\alpha) > \omega$, then, for any $U \subseteq \Omega$, U contains a neighborhood of α iff $\alpha \in U$ and U contains a club in α .

As an immediate corollary of Proposition 11.3 we obtain

Theorem 11.5. τ_2 coincides with the club topology.

We remark that the above theorem saves us the little work of verifying that the club topology is, indeed, a topology.

The derivative operation for the club topology is also well-known in set theory. Recall the following definition for $\text{cf}(\alpha) > \omega$.

A subset $A \subseteq \alpha$ is called *stationary in α* if A intersects every club in α . Observe that this happens exactly when α is a limit point of A in τ_2 , so

$$d_2(A) = \{\alpha : \text{cf}(\alpha) > \omega \text{ and } A \cap \alpha \text{ is stationary in } \alpha\}.$$

The map d_2 is usually called the *Mahlo operation* (see [36] where d_2 is denoted Tr). Its main significance is associated with the notion of Mahlo cardinal, one of the basic examples of large cardinals in set theory. Let Reg denote the class of regular cardinals; the ordinals in $d_2(\text{Reg})$ are called *weakly Mahlo cardinals*. Their existence implies the consistency of ZFC, as well as the consistency of ZFC together with the assertion ‘inaccessible cardinals exist.’

Now we turn to topology τ_3 .

11.2 Stationary reflection and Mahlo topology

Since the open sets of τ_3 are generated by the Mahlo operation, we call τ_3 *Mahlo topology*. It turns out to be intrinsically connected with the concept of stationary reflection, an extensively studied concept in set theory.

We adopt the following terminology. An ordinal λ is called *reflecting* if $\text{cf}(\lambda) > \omega$ and, whenever A is stationary in λ , there is an $\alpha < \lambda$ such that $A \cap \alpha$ is stationary in α . Similarly, λ is *doubly reflecting* if $\text{cf}(\lambda) > \omega$ and whenever A, B are stationary in λ there is an $\alpha < \lambda$ such that both $A \cap \alpha$ and $B \cap \alpha$ are stationary in α .

Mekler and Shelah’s notion of *reflection cardinal* [43] is somewhat more general than the one given here, however it has the same consistency strength. Reflection for pairs of stationary sets has been introduced by Magidor [41]. Since d_2 coincides with the Mahlo operation, we immediately obtain the following statement.

Proposition 11.6. (i) λ is reflecting iff λ is d_2 -reflexive;

(ii) λ is doubly reflecting iff λ is doubly d_2 -reflexive;

(iii) λ is a non-isolated point in τ_3 iff λ is doubly reflecting.

Together with the next proposition this yields a characterization of Mahlo topology in terms of neighborhoods.

Proposition 11.7. Suppose λ is doubly reflecting. For any subset $U \subseteq \Omega$, the following conditions are equivalent:

(i) U contains a τ_3 -neighborhood of λ ;

(ii) $\lambda \in U$ and there is a $B \subseteq \lambda$ such that $\lambda \in d_2 B \subseteq U$;

(iii) $\lambda \in U$ and there is a τ_2 -closed (in the relative topology of λ) stationary $C \subseteq \lambda$ such that $C \subseteq U$.

Notice that the notion of τ_2 -closed stationary C in (iii) is the analog of the notion of club for the τ_2 -topology.

Proof. Condition (ii) implies (iii). Since λ is reflecting, if $\lambda \in d_2B$ then $\lambda \in d_2d_2B$, that is, $\lambda \cap d_2B$ is stationary in λ . So we can take $C := \lambda \cap d_2B$.

Condition (iii) implies (ii). If C is τ_2 -closed and stationary in λ , then $d_2C \subseteq C \cup \{\lambda\} \subseteq U$ and $\lambda \in d_2C$. Thus, $\lambda \cap d_2C$ can be taken for B .

Condition (ii) implies (i). If (ii) holds, U contains a subset of the form d_2B . The latter is τ_3 -open and contains λ , thus, is a neighborhood of λ .

For the converse direction, we note that by Proposition 10.4 U contains a subset of the form $A \cap d_2B$ where $A \in \tau_2$, $B \subseteq A$ and $\lambda \in A \cap d_2B$. Since A is a τ_2 -neighborhood of λ , by Theorem 11.5 there is a set B_1 such that $\lambda \in [0, \lambda] \cap d_1B_1 \subseteq A$. Then,

$$\lambda \in [0, \lambda] \cap d_1B_1 \cap d_2B = [0, \lambda] \cap d_2(B \cap d_1B_1).$$

Since $[0, \lambda]$ is clopen, we obtain $\lambda \in d_2C$ with $C := B \cap d_1B_1 \cap \lambda$. ⊣

Reflecting and doubly reflecting cardinals are large cardinals in the sense that their existence implies consistency of ZFC. They have been studied by Mekler and Shelah [43] and Magidor [41] who investigated their consistency strength and related them to some other well-known large cardinals. By a result of Magidor, the existence of a doubly reflecting cardinal is equiconsistent with the existence of a *weakly compact cardinal*.⁵ More precisely, the following proposition holds.

Proposition 11.8. (i) *If λ is weakly compact then λ is doubly reflecting.*

(ii) (Magidor) *If λ is doubly reflecting then λ is weakly compact in L .*

Here, the first item is well-known and easy. Magidor originally proved the analog of the second item for $\lambda = \aleph_2$ and stationary sets of ordinals of countable cofinality in \aleph_2 . However, it has been remarked by Mekler and Shelah [43] that essentially the same proof yields the stated claim.⁶

Corollary 11.9. *Assertion “ τ_3 is non-discrete” is equiconsistent with the existence of a weakly compact cardinal.*

Corollary 11.10. *If ZFC is consistent then it is consistent with ZFC that τ_3 is discrete and hence that \mathbf{GLP}_3 is incomplete w.r.t. any ordinal space.*

Recall that θ_n denotes the first non-isolated point of τ_n (in the space of all ordinals). We have: $\theta_0 = 1$, $\theta_1 = \omega$, $\theta_2 = \omega_1$, θ_3 is the first doubly reflecting cardinal.

ZFC does not know much about the location of θ_3 , however the following facts are interesting.

- θ_3 is regular, but not a successor of a regular cardinal;
- While weakly compact cardinals are non-isolated, θ_3 need not be weakly compact: If infinitely many supercompact cardinals exist, then there is a model where $\aleph_{\omega+1}$ is doubly reflecting (Magidor [41]);
- If θ_3 is a successor of a singular cardinal, then some very strong large cardinal hypothesis must be consistent (Woodin cardinals).

⁵Weakly compact cardinals are the same as Π_1^1 -indescribable cardinals, see below.

⁶The first author thanks J. Cummings for clarifying this.

11.3 Further topologies

Further topologies of the ordinal GLP-space do not seem to have prominently occurred in set-theoretic work. They yield some large cardinal notions, for the statement that τ_n is non-discrete (or, equivalently, the statement that θ_n exists) implies the existence of a doubly reflecting cardinal, for any $n > 2$. We do not know whether cardinals θ_n coincide with any of the standard large cardinal notions.

Here we give a sufficient condition for the topology τ_{n+2} to be non-discrete. We show that, if there exists a Π_n^1 -indescribable cardinal, then τ_{n+2} is non-discrete.

Let Q be a class of second order formulas over the standard first order set-theoretic language enriched by a unary predicate R . We assume Q to contain at least the class of all first order formulas (denoted Π_0^1). We shall consider standard models of that language of the form (V_α, \in, R) , where α is an ordinal, V_α is the α -th class in the cumulative hierarchy, and R is a subset of V_α .

We would like to give a definition of Q -indescribable cardinals in topological terms. They can then be defined as follows.

Definition 11.11. For any sentence $\varphi \in Q$ and any $R \subseteq V_\kappa$, let $U_\kappa(\varphi, R)$ denote the set

$$\{\alpha \leq \kappa : (V_\alpha, \in, R \cap V_\alpha) \models \varphi\}.$$

The Q -describable topology τ_Q on Ω is generated by a subbase consisting of sets $U_\kappa(\varphi, R)$, for all $\kappa \in \Omega$, $\varphi \in Q$, and all $R \subseteq V_\kappa$.

As an exercise, the reader can check that the intervals $(\alpha, \beta]$ are open in any τ_Q (consider $R = \{\alpha\}$ and $\varphi = \exists x (x \in R)$). The main strength of the Q -describable topology, however, comes from the fact that a second order variable R is allowed to occur in φ . So, a lot of subsets of Ω that can be ‘described’ in this way are open in τ_Q .

Let d_Q denote the derivative operator for τ_Q . An ordinal κ is called Q -indescribable if it is a limit point of τ_Q . In other words, κ is Q -indescribable iff $\kappa \in d_Q(\text{On})$ iff $\kappa \in d_Q(\kappa)$.

It is not difficult to show that, whenever Q is any of the classes Π_n^1 , the sets $U_\kappa(\varphi, R)$ actually form a base for τ_Q . Hence, our definition of Π_n^1 -indescribable cardinals is equivalent to the standard one given in [37]: κ is Q -indescribable iff, for all $R \subseteq V_\kappa$ and all sentences $\varphi \in Q$,

$$(V_\kappa, \in, R) \models \varphi \Rightarrow \exists \alpha < \kappa (V_\alpha, \in, R \cap V_\alpha) \models \varphi.$$

A *weakly compact cardinal* can be defined as the Π_1^1 -indescribable one. The well-known fact that weakly compact cardinals are doubly reflecting can be somewhat more generally stated as follows: the Mahlo topology τ_3 is contained in $\tau_{\Pi_1^1}$. We omit a short proof, because we are going to prove a more general proposition suggested to the first author by Philipp Schlicht (see [10]).

Proposition 11.12. For any $n \geq 0$, τ_{n+2} is contained in $\tau_{\Pi_n^1}$.

Proof. We shall show that, for each n , there is a Π_n^1 -formula $\varphi_{n+1}(R)$ such that

$$\kappa \in d_{n+1}(A) \iff (V_\kappa, \in, A \cap \kappa) \models \varphi_{n+1}(R). \quad (**)$$

This implies that, for each $\kappa \in d_{n+1}(A)$, the set $U_\kappa(\varphi_{n+1}, A \cap \kappa)$ is a $\tau_{\Pi_n^1}$ -open subset of $d_{n+1}(A)$ containing κ . Hence, each $d_{n+1}(A)$ is $\tau_{\Pi_n^1}$ -open. Since τ_{n+2} is

generated over τ_{n+1} by the open sets of the form $d_{n+1}(A)$ for various A , we have $\tau_{n+2} \subseteq \tau_{\Pi_n^1}$.

We prove $(**)$ by induction on n . For $n = 0$, notice that $\kappa \in d_1(A)$ iff $(\kappa \in \text{Lim and } A \cap \kappa \text{ is unbounded in } \kappa)$ iff

$$(V_\kappa, \in, A \cap \kappa) \models \forall \alpha \exists \beta (R(\beta) \wedge \alpha < \beta).$$

For the induction step recall that, by Corollary 10.5, $\kappa \in d_{n+1}(A)$ iff

- (i) κ is doubly d_n -reflexive;
- (ii) $\forall Y \subseteq \kappa (\kappa \in d_n(Y) \rightarrow \exists \alpha < \kappa (\alpha \in A \wedge \alpha \in d_n(Y)))$.

By the induction hypothesis, for some $\varphi_n(R) \in \Pi_{n-1}^1$, we have

$$\alpha \in d_n(A) \iff (V_\alpha, \in, A \cap \alpha) \models \varphi_n(R).$$

Hence, part (ii) is equivalent to

$$(V_\kappa, \in, A \cap \kappa) \models \forall Y \subseteq \text{On} (\varphi_n(Y) \rightarrow \exists \alpha (R(\alpha) \wedge \varphi_n^{V_\alpha}(Y \cap \alpha))).$$

Here, φ^{V_α} means the relativization of all quantifiers in φ to V_α . We notice that V_α is first order definable, hence the complexity of $\varphi_n^{V_\alpha}$ remains in the class Π_{n-1}^1 . So, the resulting formula is Π_n^1 .

To treat part (i) we recall that κ is doubly d_n -reflexive iff $\kappa \in d_n(\text{On})$ and

$$\forall Y_1, Y_2 \subseteq \kappa (\kappa \in d_n(Y_1) \cap d_n(Y_2) \rightarrow \exists \alpha < \kappa \alpha \in d_n(Y_1) \cap d_n(Y_2)).$$

Similarly to the above, using the induction hypothesis this can be rewritten as a Π_n^1 -formula. \dashv

Corollary 11.13. *If there is a Π_n^1 -indescribable cardinal, then τ_{n+2} has a non-isolated point.*

Corollary 11.14. *If there is a cardinal which is Π_n^1 -indescribable, for each n , then all τ_n are non-discrete.*

We know that θ_3 need not be weakly compact in some models of ZFC. Hence, the condition of the existence of Π_n^1 -indescribable cardinals is not a necessary one for the nontriviality of the topologies τ_{n+2} , in general. However, the two statements are equiconsistent, as has been recently shown by Bagaria, Magidor, and Sakai in [4].

12 Topological completeness results

As in the case of the unimodal language (cf Section 3), one can ask two basic questions:

- Is **GLP** complete w.r.t. the class of all GLP-spaces?
- Is **GLP** complete w.r.t. some fixed natural GLP-spaces?

In the unimodal case, both questions received positive answers due to Esakia and Abashidze–Blass, respectively. Now the situation is more complicated.

12.1 Topological completeness of GLP

The first question was initially studied by Beklemishev, Bezhanishvili and Icard in [14]. In this paper only some partial results were obtained. It was proved that the bimodal system **GLP**₂ is complete w.r.t. GLP₂-spaces of the form (X, τ, τ^+) , where X is a well-founded partial ordering and τ is its left topology. A proof of this result was based on the Kripke model techniques coming from [11].

Already at that time it was clear that these techniques cannot be immediately generalized to GLP₃-spaces, since the third topology τ^{++} on such orderings is sufficiently similar to the club topology. From the results of Andreas Blass described below it was known that some stronger set-theoretic assumptions would be needed to prove the completeness w.r.t. such topologies. Moreover, without any large cardinal assumptions it was not even known whether a GLP-space with a non-discrete third topology could exist at all.

First examples of GLP-spaces in which all topologies are non-discrete are constructed in [5]. In the same paper also the much stronger fact of topological completeness of **GLP** w.r.t. the class of all (countable, Hausdorff) GLP-spaces is established.

Theorem 12.1. (i) $\text{Log}(\mathcal{C}) = \mathbf{GLP}$, where \mathcal{C} is the class of all GLP-spaces.
(ii) There is a countable Hausdorff GLP-space X such that $\text{Log}(X) = \mathbf{GLP}$.

In fact, X is the ordinal ε_0 equipped with a sequence of topologies refining the order topology. However, these topologies cannot be first-countable and are, in fact, defined using non-constructive methods such as Zorn's lemma. In this sense, it is not an example of a *natural* GLP-space.

A proof of this theorem introduces the techniques of maximal rank-preserving and limit-maximal extensions of scattered spaces. It falls outside the present survey (see [5]).

12.2 Completeness w.r.t. the ordinal GLP-space

The question whether **GLP** is complete w.r.t. some natural GLP-space is still open. Some partial results concerning the GLP-space generated from the order topology on the ordinals are described below. Here, we call this space the *ordinal GLP-space*. (The space described in the previous section is clearly not an exact model of **GLP**, as the left topology validates the linearity axiom.)

As we know from Corollary 11.10, it is consistent with ZFC that the Mahlo topology is discrete. Hence, it is consistent that **GLP** is incomplete w.r.t. the ordinal GLP-space. However, is it consistent that **GLP** be complete? To this question we do not know a full answer. A pioneering work has been done by Andreas Blass [18] who studied the question of completeness of the Gödel–Löb logic **GL** w.r.t. a semantics equivalent to the topological interpretation w.r.t. the *club topology* τ_2 . He used the language of filters rather than that of topological spaces, as is more common in set theory.

Theorem 12.2 (Blass). (i) If $V = L$ and $\Omega \geq \aleph_\omega$, then **GL** is complete w.r.t. (Ω, τ_2) .
(ii) If there is a weakly Mahlo cardinal, there is a model of ZFC in which **GL** is incomplete w.r.t. (Ω, τ_2) , for any Ω .

A corollary of (i) is that the statement “**GL** is complete w.r.t. τ_2 ” is consistent with ZFC (provided ZFC is consistent). In fact, instead of $V = L$ Blass used the so-called *square principle* for all \aleph_n , $n < \omega$, which holds in L by the results of Ronald Jensen. A proof of (i) is based on an interesting combinatorial construction using the techniques of splitting stationary sets.

A proof of (ii) is much easier. It uses a model of Harrington and Shelah in which \aleph_2 is reflecting for stationary sets of ordinals of countable cofinality [31]. Assuming Mahlo cardinals exist, they have shown that the following statement holds in some model of ZFC:

If S is a stationary subset of \aleph_2 such that $\forall \alpha \in S \text{ cf}(\alpha) = \omega$, then there is a $\beta < \alpha$ (of cofinality ω_1) such that $S \cap \beta$ is stationary in β .

In fact, this statement can be expressed in the language of modal logic. First, we remark that this principle implies its generalization to all ordinals λ of cofinality \aleph_2 (consider an increasing continuous function mapping \aleph_2 to a club in λ). Second, we remark that, for the club topology, the formula $\Diamond^n \top$ represents the class ordinals of cofinality at least \aleph_n . This is a straightforward generalization of Lemma 11.1. Thus, the formula $\Box^3 \perp \wedge \Diamond^2 \top$ represents the subclass of Ω consisting of ordinals of cofinality ω_2 .

Hence, the above reflection principle amounts to the validity of the following modal formula:

$$\Box^3 \perp \wedge \Diamond^2 \top \wedge \Diamond(p \wedge \Box \perp) \rightarrow \Diamond^2(p \wedge \Box \perp). \quad (*)$$

In fact, if the left hand side is valid in λ , then $\text{cf}(\lambda) = \omega_2$ and the interpretation of $p \wedge \Box \perp$ is a set S consisting of ordinals of countable cofinality such that $S \cap \lambda$ is stationary in λ . The right hand side just states that this set reflects. Thus, formula $(*)$ is valid in (Ω, τ_2) , for any Ω . Since this formula is clearly not provable in **GL**, the topological completeness fails for (Ω, τ_2) .

Thus, Blass managed to give an exact consistency strength of the statement “**GL** is incomplete w.r.t. τ_2 ”.

Corollary 12.3. “**GL** is incomplete w.r.t. τ_2 ” is consistent iff it is consistent that Mahlo cardinals exist.

It is possible to generalize these results to the case of bimodal logic **GLP**₂ [12]. The situation remains essentially unchanged, although a proof of Statement (i) of Blass’s theorem needs considerable adaptation.

Theorem 12.4. If $V = L$ and $\Omega \geq \aleph_\omega$, then **GLP**₂ is complete w.r.t. $(\Omega; \tau_1, \tau_2)$.

13 Topologies for the variable-free fragment of GLP

A natural topological model for the variable-free fragment of **GLP** has been introduced by Thomas Icard [33]. It is not a GLP-space, thus, not a model of the full **GLP**. However, it is sound and complete for the variable-free fragment of **GLP**. It gives a convenient tool for the study of this fragment, which plays an important role in proof-theoretic applications of the polymodal provability logic. Here, we give a simplified presentation of Icard’s polytopological space.

Let Ω be an ordinal and let $\ell : \Omega \rightarrow \Omega$ denote the rank function for the order topology on Ω . It is easy to check that $\ell(0) = 0$ and $\ell(\alpha) = \beta$ whenever $\alpha = \gamma + \omega^\beta$. By the Cantor normal form theorem, such a β is uniquely defined for each ordinal $\alpha > 0$. We define $\ell^0(\alpha) = \alpha$ and $\ell^{k+1}(\alpha) = \ell\ell^k(\alpha)$.

Icard's topologies v_n , for each $n \in \omega$, are defined as follows. Let v_0 be the left topology, and let v_n be generated by v_0 and all sets of the form

$$U_\beta^m := \{\alpha \in \Omega : \ell^m(\alpha) > \beta\},$$

for $m < n$.

Clearly, v_n is an increasing sequence of topologies. In fact, v_1 is the standard interval topology. We let d_n and ρ_n denote the derivative operator and the rank function for v_n , respectively. We have the following characterizations.

- Lemma 13.1.** (i) $\ell : (\Omega, v_{n+1}) \rightarrow (\Omega, v_n)$ is a d -map;
(ii) v_{n+1} is the coarsest topology ν on Ω such that ν contains the interval topology and $\ell : (\Omega, \nu) \rightarrow (\Omega, v_n)$ is continuous;
(iii) ℓ^n is the rank function of v_n , that is, $\rho_n = \ell^n$;
(iv) v_{n+1} is generated by v_n and $\{d_n^{\alpha+1}(\Omega) : \alpha < \rho_n(\Omega)\}$.

Proof. (i) The map $\ell : (\Omega, v_{n+1}) \rightarrow (\Omega, v_n)$ is continuous. In fact, $\ell^{-1}[0, \beta)$ is open in the order topology v_1 , since $\ell : (\Omega, v_1) \rightarrow (\Omega, v_0)$ is its rank function, hence a d -map. Also, if $m < n$ then $\ell^{-1}(U_\beta^m) = U_\beta^{m+1}$, hence it is open in v_{n+1} .

The map ℓ is open. Notice that v_{n+1} is generated by v_1 and some sets of the form $\ell^{-1}(U)$ where $U \in v_n$. A base of v_{n+1} consists of sets of the form $V \cap \ell^{-1}(U)$, for some $V \in v_1$ and $U \in v_n$. We have $\ell(V \cap \ell^{-1}(U)) = \ell(V) \cap U$. $\ell(V)$ is v_0 -open, since $\ell : (\Omega, v_1) \rightarrow (\Omega, v_0)$ is a d -map and $V \in v_1$. Hence, the image of any basic open in v_{n+1} is open in v_n .

The map ℓ is pointwise discrete, since $\ell^{-1}\{\alpha\}$ is discrete in the order topology v_1 , hence in v_{n+1} .

(ii) By (i), $\ell : (\Omega, v_{n+1}) \rightarrow (\Omega, v_n)$ is continuous, hence $\nu \subseteq v_{n+1}$. On the other hand, if $\ell : (\Omega, \nu) \rightarrow (\Omega, v_n)$ is continuous, then $\ell^{-1}(U_\beta^m) \in \nu$, for each $m < n$. Therefore $U_\beta^m \in \nu$, for all m such that $1 \leq m \leq n$. Since ν also contains the interval topology, we have $v_{n+1} \subseteq \nu$.

(iii) By (i), we have that $\rho_n \circ \ell$ is a d -map from (Ω, v_{n+1}) to (Ω, v_0) . Hence, it coincides with the rank function for v_{n+1} , $\rho_{n+1} = \rho_n \circ \ell$. The claim follows by an easy induction on n .

(iv) Notice that, by (iii),

$$d_n^{\beta+1}(\Omega) = \{\alpha \in \Omega : \rho_n(\alpha) > \beta\} = \{\alpha \in \Omega : \ell^n(\alpha) > \beta\} = U_\beta^n.$$

Obviously, v_{n+1} is generated by v_n and U_β^n , for all β . Hence, the claim. \dashv

We call *Icard space* a polytopological space of the form $(\Omega; v_0, v_1, \dots)$. Icard originally considered just $\Omega = \varepsilon_0$. We are going to give an alternative proof of the following theorem [33].

Theorem 13.2 (Icard). *Let φ be a variable-free GLP-formula.*

- (i) *If $\mathbf{GLP} \vdash \varphi$, then $(\Omega; v_0, v_1, \dots) \models \varphi$.*
- (ii) *If $\Omega \geq \varepsilon_0$ and $\mathbf{GLP} \not\vdash \varphi$, then $(\Omega; v_0, v_1, \dots) \not\models \varphi$.*

Proof. Within this proof we abbreviate $(\Omega; v_0, v_1, \dots)$ by Ω . To prove part (i) we first remark that all topologies v_n are scattered, hence all axioms of **GLP** except for P1 are valid in Ω . Moreover, $\text{Log}(\Omega)$ is closed under the inference rules of **GLP**. Thus, we only have to show that the variable-free instances of axiom P1 are valid in Ω . This is sufficient, because any derivation of a variable-free formula in **GLP** can be replaced by a derivation in which only the variable-free formulas occur (replace all the variables by the constant \top).

Let φ be a variable-free formula. We denote by φ^* its uniquely defined interpretation in Ω . The validity of an instance of P1 for φ amounts to the fact that $d_m(\varphi^*)$ is open in v_n whenever $m < n$. Thus, we have to prove the following proposition.

Proposition 13.3. *For any variable-free formula φ , $d_n(\varphi^*)$ is open in v_{n+1} .*

Let φ^+ denote the result of replacing in φ each modality $\langle n \rangle$ by $\langle n+1 \rangle$. We need the following auxiliary claim.

Lemma 13.4. *If φ is variable-free, then $\ell^{-1}(\varphi^*) = (\varphi^+)^*$.*

Proof. This goes by induction on the build-up of φ . The cases of constants and boolean connectives are easy. Suppose $\varphi = \langle n \rangle \psi$. We notice that since $\ell : (\Omega, v_{n+1}) \rightarrow (\Omega, v_n)$ is a d -map, we have $\ell^{-1}(d_n(A)) = d_{n+1}(\ell^{-1}(A))$, for any $A \subseteq \Omega$. Therefore,

$$\ell^{-1}(\varphi^*) = \ell^{-1}(d_n(\psi^*)) = d_{n+1}(\ell^{-1}(\psi^*)) = d_{n+1}((\psi^+)^*) = (\varphi^+)^*,$$

as required. \dashv

We prove Proposition 13.3 in two steps. First, we show that it holds for a subclass of variable-free formulas called *ordered formulas*. Then we show that any variable-free formula is equivalent in Ω to an ordered one.

A formula φ is called *ordered*, if no modality $\langle m \rangle$ occurs within the scope of $\langle n \rangle$ in φ for any $m < n$. The *height* of φ is the index of its maximal modality.

Lemma 13.5. *If $\langle n \rangle \varphi$ is ordered, then $d_n(\varphi^*)$ is open in v_{n+1} .*

Proof. This goes by induction on the height of $\langle n \rangle \varphi$. If it is 0, then $n = 0$. If $n = 0$, the claim is obvious, since $d_0(A)$ is open in v_1 , for any $A \subseteq \Omega$. If $n > 0$, since $\langle n \rangle \varphi$ is ordered, we observe that $\langle n \rangle \varphi$ has the form $(\langle n-1 \rangle \psi)^+$, for some ψ . The height of $\langle n-1 \rangle \psi$ is less than that of $\langle n \rangle \varphi$. Hence, by the induction hypothesis, $(\langle n-1 \rangle \psi)^* \in v_n$. Since $\ell : (\Omega, v_{n+1}) \rightarrow (\Omega, v_n)$ is continuous, we conclude that $\ell^{-1}(\langle n-1 \rangle \psi)^*$ is open in v_{n+1} . By Lemma 13.4, this set coincides with $(\langle n \rangle \varphi)^* = d_n(\varphi^*)$. \dashv

Lemma 13.6. *Any variable-free formula φ is equivalent in Ω to an ordered one.*

Proof. We argue by induction on the complexity of φ . The cases of boolean connectives and constants are easy. Suppose φ has the form $\langle n \rangle \psi$, where we can assume ψ to be in disjunctive normal form $\psi = \bigvee_i \bigwedge_j \pm \langle n_{ij} \rangle \psi_{ij}$. By the induction hypothesis, we may assume all the subformulas $\langle n_{ij} \rangle \psi_{ij}$ (and ψ itself) ordered. Since $\langle n \rangle$ commutes with disjunction, it will be sufficient to show that, for each i , the formula

$$\langle n \rangle \bigwedge_j \pm \langle n_{ij} \rangle \psi_{ij} \tag{*}$$

can be ordered.

By Lemma 13.5 each set $(\langle n_{ij} \rangle \psi_{ij})^*$ is open in v_n , whenever $n_{ij} < n$. Being a derived set, it is also closed in $v_{n_{ij}}$ and hence in v_n . Thus, all such sets are clopen.

If U is open then $d(A \cap U) = d(A) \cap U$, for any topological space. In particular, for any $A \subseteq \Omega$ and $n_{ij} < n$,

$$d_n(A \cap (\pm \langle n_{ij} \rangle \psi_{ij})^*) = d_n(A) \cap (\pm \langle n_{ij} \rangle \psi_{ij})^*.$$

This allows us to bring all the conjuncts $\pm \langle n_{ij} \rangle \psi_{ij}$ from under the $\langle n \rangle$ modality in $(*)$. The resulting conjunction is ordered. \dashv

This concludes the proof of Proposition 13.3 and thereby of Part (i).

To prove Part (ii), we shall rely on the following fundamental lemma about the variable-free fragment of **GLP**. For a purely syntactic proof of this lemma we refer the reader to [6, 8].

A variable-free formula A is called a *word* if it is built-up from \top only using connectives of the form $\langle n \rangle$, for any $n \in \omega$. We write $A \vdash B$ for **GLP** $\vdash A \rightarrow B$.

- Lemma 13.7.** (i) *Every variable-free formula is equivalent in **GLP** to a boolean combination of words;*
(ii) *For any words A and B , either $A \vdash \langle 0 \rangle B$, or $B \vdash \langle 0 \rangle A$, or A and B are equivalent;*
(iii) *Conjunction of words is equivalent to a word.*

We prove Part (ii) of Theorem 13.2 in a series of lemmas. First, we show that any word is satisfiable in Ω provided $\Omega \geq \varepsilon_0$.

Lemma 13.8. *For any word A , $\varepsilon_0 \in A^*$.*

Proof. We know that $\rho_n(\varepsilon_0) = \ell^n(\varepsilon_0) = \varepsilon_0$. Hence, $\varepsilon_0 \in d_n(\Omega)$, for each n . Assume n exceeds all the indices of modalities in A and $A = \langle m \rangle B$. By Proposition 13.3 the set B^* is open in v_n . By the induction hypothesis $\varepsilon_0 \in B^*$. Hence, $\varepsilon_0 \in d_n(B^*) \subseteq d_m(B^*) = A^*$. This proves the claim. \dashv

Applying this lemma to the word $\langle 0 \rangle A$ we obtain the following corollary.

Corollary 13.9. *For every word A , there is an $\alpha < \varepsilon_0$ such that $\alpha \in A^*$.*

Let $\min(A^*)$ denote the least ordinal $\alpha \in \Omega$ such that $\alpha \in A^*$.

Lemma 13.10. *For any words A, B , if $A \not\vdash B$ then $\min(A^*) \notin B^*$.*

Proof. If $A \not\vdash B$ then, by Lemma 13.7 (ii), $B \vdash \langle 0 \rangle A$. Therefore, by the soundness of **GLP** in Ω , $B^* \subseteq d_0(A^*)$. It follows that for each $\beta \in B^*$ there is an $\alpha \in A^*$ such that $\alpha < \beta$. Thus, $\min(A^*) \notin B^*$. \dashv

Now we are ready to prove Part (ii). Assume φ is variable-free and **GLP** $\not\vdash \varphi$. By Lemma 13.7 (i) we can assume that φ is a boolean combination of words. Writing φ in conjunctive normal form we observe that it is sufficient to prove the claim only for formulas φ of the form $\bigwedge_i A_i \rightarrow \bigvee_j B_j$, where A_i and B_j are words. Moreover, $\bigwedge_i A_i$ is equivalent to a single word A .

Since **GLP** $\not\vdash \varphi$ we have $A \not\vdash B_j$, for each j . Let $\alpha = \min(A^*)$. By Lemma 13.10 we have $\alpha \notin B_j^*$, for each j . Hence, $\alpha \notin (\bigvee_j B_j)^*$ and $\alpha \notin \varphi^*$. This means that $\Omega \not\vdash \varphi^*$. \dashv

14 Further results

Topological semantics of polymodal provability logic has been extended to the language with transfinitely many modalities. A logic \mathbf{GLP}_Λ having modalities $[\alpha]$, for all ordinals $\alpha < \Lambda$, is introduced in [8]. It was intended for the proof-theoretic analysis of predicative theories and is currently being actively investigated for that purpose.

David Fernandez and Joost Joosten undertook a thorough study of the variable-free fragment of that logic mostly in connection with the arising ordinal notation systems (see [24, 26] for a sample). In particular, they found a suitable generalization of Icard’s polytopological space and showed that it is complete for that fragment [25]. Fernandez [29] also proved topological completeness of the full \mathbf{GLP}_Λ by generalizing the results in [5].

The ordinal GLP-space is easily generalized to transfinitely many topologies $(\tau_\alpha)_{\alpha < \Lambda}$ by letting τ_0 be the left topology, $\tau_{\alpha+1} := \tau_\alpha^+$ and, for limit ordinals λ , τ_λ be the topology generated by all τ_α such that $\alpha < \lambda$. This space is a natural model of \mathbf{GLP}_Λ and has been studied quite recently by Joan Bagaria [3] and further by Bagaria, Magidor and Sakai [4]. In particular, the three authors proved that in L the limit points of τ_{n+2} are Π_n^1 -indescribable cardinals. The question posed in [14] whether the non-discreteness of τ_{n+2} is equiconsistent with the existence of Π_n^1 -indescribable cardinals still seems to be open.

15 Acknowledgement

The first author was supported by the Russian Foundation for Basic Research (RFBR), Russian Presidential Council for Support of Leading Scientific Schools, and the Swiss–Russian cooperation project STCP–CH–RU “Computational proof theory”.

The second author was supported by the Shota Rustaveli National Science Foundation grant #FR/489/5-105/11 and the French–Georgian grant CNRS–SRNSF #4135/05-01.

References

- [1] M. Abashidze. Ordinal completeness of the Gödel–Löb modal system. In *Intensional logics and the logical structure of theories*, pages 49–73. Metsniereba, Tbilisi, 1985. In Russian.
- [2] S.N. Artemov and L.D. Beklemishev. Provability logic. In D. Gabbay and F. Guenther, editors, *Handbook of Philosophical Logic, 2nd ed.*, volume 13, pages 229–403. Kluwer, Dordrecht, 2004.
- [3] J. Bagaria. Topologies on ordinals and the completeness of polymodal provability logic. In preparation, 2012.
- [4] J. Bagaria, M. Magidor, and H. Sakai. Reflection and indescribability in the constructible universe. Manuscript, 2012.
- [5] L. Beklemishev and D. Gabelaia. Topological completeness of the provability logic GLP. Preprint arXiv:1106.5693v1 [math.LO], 2011.

- [6] L.D. Beklemishev. Provability algebras and proof-theoretic ordinals, I. *Annals of Pure and Applied Logic*, 128:103–123, 2004.
- [7] L.D. Beklemishev. Reflection principles and provability algebras in formal arithmetic. *Uspekhi Matematicheskikh Nauk*, 60(2):3–78, 2005. In Russian. English translation in: *Russian Mathematical Surveys*, 60(2): 197–268, 2005.
- [8] L.D. Beklemishev. Veblen hierarchy in the context of provability algebras. In P. Hájek, L. Valdés-Villanueva, and D. Westerståhl, editors, *Logic, Methodology and Philosophy of Science, Proceedings of the Twelfth International Congress*, pages 65–78. Kings College Publications, London, 2005. Preprint: Logic Group Preprint Series 232, Utrecht University, June 2004.
- [9] L.D. Beklemishev. The Worm Principle. In Z. Chatzidakis, P. Koepke, and W. Pohlers, editors, *Lecture Notes in Logic 27. Logic Colloquium '02*, pages 75–95. AK Peters, 2006. Preprint: Logic Group Preprint Series 219, Utrecht University, March 2003.
- [10] L.D. Beklemishev. On GLP-spaces. Manuscript, [http://www.mi.ras.ru/~ bekl/Papers/glp-sp.pdf](http://www.mi.ras.ru/~bekl/Papers/glp-sp.pdf), 2009.
- [11] L.D. Beklemishev. Kripke semantics for provability logic GLP. *Annals of Pure and Applied Logic*, 161:756–774, 2010. Preprint: Logic Group Preprint Series 260, University of Utrecht, November 2007. <http://preprints.phil.uu.nl/lgps/>.
- [12] L.D. Beklemishev. Ordinal completeness of bimodal provability logic GLB. In N. Bezhanishvili et al., editor, *Logic, Language and Computation, TbiLLC 2009. Lecture Notes in Artificial Intelligence*, number 6618, pages 1–15, 2011.
- [13] L.D. Beklemishev. A simplified proof of the arithmetical completeness theorem for the provability logic GLP. *Trudy Matematicheskogo Instituta imeni V.A.Steklova*, 274(3):32–40, 2011. In Russian. English translation: *Proceedings of the Steklov Institute of Mathematics*, 274(3):25–33, 2011.
- [14] L.D. Beklemishev, G. Bezhanishvili, and T. Icard. On topological models of GLP. In R. Schindler, editor, *Ways of Proof Theory*, Ontos Mathematical Logic, pages 133–152. Ontos Verlag, Heusendamm bei Frankfurt, Germany, 2010. Preprint: Logic Group Preprint Series 278, University of Utrecht, August 2009, <http://preprints.phil.uu.nl/lgps/>.
- [15] L.D. Beklemishev, J. Joosten, and M. Vervoort. A finitary treatment of the closed fragment of Japaridze’s provability logic. *Journal of Logic and Computation*, 15(4):447–463, 2005.
- [16] G. Bezhanishvili, L. Esakia, and D. Gabelaia. Some results on modal axiomatization and definability for topological spaces. *Studia Logica*, 81:325–355, 2005.
- [17] G. Bezhanishvili and P. Morandi. Scattered and hereditarily irresolvable spaces in modal logic. *Archive*, 49(3):343–365, 2010.
- [18] A. Blass. Infinitary combinatorics and modal logic. *Journal of Symbolic Logic*, 55(2):761–778, 1990.
- [19] G. Boolos. *The Unprovability of Consistency: An Essay in Modal Logic*. Cambridge University Press, Cambridge, 1979.
- [20] G. Boolos. The analytical completeness of Dzjaparidze’s polymodal logics. *Annals of Pure and Applied Logic*, 61:95–111, 1993.

- [21] G. Boolos. *The Logic of Provability*. Cambridge University Press, Cambridge, 1993.
- [22] L. Carlucci. Worms, gaps and hydras. *Mathematical Logic Quarterly*, 51(4):342–350, 2005.
- [23] A. Chagrov and M. Zakharyashev. *Modal Logic*. Clarendon Press, Oxford, 1997.
- [24] D. Fernández-Duque and J. J. Joosten. Kripke models of transfinite provability logic. In *Advances in Modal Logic*, volume 9, pages 185–199. College Publications, 2012.
- [25] D. Fernández-Duque and J. J. Joosten. Models of transfinite provability logic. To appear in *Journal of Symbolic Logic*. Preliminary version at <http://personal.us.es/dfduque/glpmodels.pdf>, 2012.
- [26] D. Fernández-Duque and J. J. Joosten. Turing progressions and their well-orders. In *How the world computes*, Lecture Notes in Computer Science, pages 212–221. Springer, 2012.
- [27] L. Esakia. Diagonal constructions, Löb’s formula and Cantor’s scattered spaces. In *Studies in logic and semantics*, pages 128–143. Metsniereba, Tbilisi, 1981. In Russian.
- [28] S. Feferman, J.R. Dawson, S.C. Kleene, G.H. Moore, R.M. Solovay, and J. van Heijenoort, editors. *Kurt Gödel Collected Works, Volume 1: Publications 1929–1936*. Oxford University Press, 1996.
- [29] D. Fernández-Duque. The polytopologies of transfinite provability logic. Preprint arXiv:1207.6595v2 [math.LO], 2012.
- [30] K. Gödel. Eine Interpretation des intuitionistischen Aussagenkalküls. *Ergebnisse Math. Kolloq.*, 4:39–40, 1933. English translation in [28], pages 301–303.
- [31] L. Harrington and S. Shelah. Some exact equiconsistency results in set theory. *Notre Dame Journal of Formal Logic*, 26:178–188, 1985.
- [32] T. Icard and J. Joosten. Provability and interpretability logics with restricted realizations. *Notre Dame Journal of Formal Logic*, 53(2):133–154, 2012.
- [33] T.F. Icard, III. A topological study of the closed fragment of GLP. *Journal of Logic and Computation*, 21(4):683–696, 2011.
- [34] K.N. Ignatiev. On strong provability predicates and the associated modal logics. *The Journal of Symbolic Logic*, 58:249–290, 1993.
- [35] G.K. Japaridze. The modal logical means of investigation of provability. Thesis in Philosophy, in Russian, Moscow, 1986.
- [36] T. Jech. *Set Theory. The Third Millennium Edition*. Springer, 2002.
- [37] A. Kanamori. *The Higher Infinite. Second edition*. Springer, 2009.
- [38] M.H. Löb. Solution of a problem of Leon Henkin. *The Journal of Symbolic Logic*, 20:115–118, 1955.
- [39] A. Macintyre and H. Simmons. Gödel’s diagonalization technique and related properties of theories. *Colloq. Math.*, 28:165–180, 1973.
- [40] R. Magari. The diagonalizable algebras (the algebraization of the theories which express Theor.:II). *Bollettino della Unione Matematica Italiana, Serie 4*, 12, 1975. Suppl. fasc. 3, 117–125.

- [41] M. Magidor. Reflecting stationary sets. *The Journal of Symbolic Logic*, 47:755–771, 1982.
- [42] J. C. C. McKinsey and A. Tarski. The algebra of topology. *Annals of Mathematics*, 45:141–191, 1944.
- [43] A. Mekler and S. Shehlah. The consistency strength of “every stationary set reflects”. *Israel Journal of Mathematics*, 67(3):353–366, 1989.
- [44] H. Simmons. Topological aspects of suitable theories. *Proc. Edinburgh Math. Soc. (2)*, 19:383–391, 1975.
- [45] C. Smoryński. *Self-Reference and Modal Logic*. Springer-Verlag, Berlin, 1985.
- [46] R.M. Solovay. Provability interpretations of modal logic. *Israel Journal of Mathematics*, 28:33–71, 1976.
- [47] C. Steinsvold. *Topological models of belief logics*. PhD thesis, New York, 2007.
- [48] A. Visser. The provability logics of recursively enumerable theories extending Peano Arithmetic at arbitrary theories extending Peano Arithmetic. *Journal of Philosophic Logic*, 13:97–113, 1984.